

FEASIBLE OBJECTIVE CONFIDENCE BOUNDS  
FOR SYSTEM RELIABILITY

BY

M. VERNON JOHNS, JR.

TECHNICAL REPORT NO. 6  
AUGUST 20, 1981

U.S. ARMY RESEARCH OFFICE  
RESEARCH TRIANGLE PARK, NORTH CAROLINA  
CONTRACT NO. DAAG29-79-C-0166

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA



FEASIBLE OBJECTIVE CONFIDENCE BOUNDS  
FOR SYSTEM RELIABILITY

BY

M. VERNON JOHNS, JR.

TECHNICAL REPORT NO. 6

AUGUST 20, 1981

U.S. ARMY RESEARCH OFFICE  
RESEARCH TRIANGLE PARK, NORTH CAROLINA  
CONTRACT NO. DAAG29-79-C-0166

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA

# FEASIBLE OBJECTIVE CONFIDENCE BOUNDS FOR SYSTEM RELIABILITY

by

M. Vernon Johns, Jr.

## ABSTRACT

The basic problem of determining objective (frequentistic) confidence bounds for the reliability of a series system based on failure data from tests of the independent components is addressed. The notion of confidence bounds based on orderings imposed on the sample space is exploited, and certain optimality considerations are incorporated. Advantage is taken of the simplifications resulting from the use of the Poisson approximation for data from highly reliable components. Tables of exact confidence bounds are produced for the case of two-component systems. These bounds are computed using sample orderings generated sequentially by a two-stage, prospective optimization procedure. A generalization of the Lindstrom-Madden technique is proposed for using the tables to find confidence bounds for systems consisting of more than two components with differing sample sizes.

Key Words: Reliability, series-system, confidence bounds, Poisson approximation, Lindstrom-Madden, sample orderings.

# FEASIBLE OBJECTIVE CONFIDENCE BOUNDS FOR SYSTEM RELIABILITY

By

M. Vernon Johns, Jr.

## 1. Introduction

Certainly one of the most basic statistical problems in the assessment of system reliability is that of determining a confidence bound on the reliability of a series system based on component failure data. The continuing appearance of papers concerned with this subject (e.g., Harris and Soms 1981, Butcher et al. 1978, and Winterbottom 1980) testifies to the elusiveness of solutions which are both practically feasible and acceptably precise. The present paper deals with the case of systems characterized by high intrinsic reliability ( $>90\%$ ) where the use of the Poisson approximation for the binomial distributions of component failure data does not introduce appreciable error. The emphasis is on objective, frequentistic confidence bounds which avoid the uncertainties of interpretation associated with posterior bounds obtained by Bayesian methods. The use of the Poisson distribution provides several advantages. It easily leads to valid results for cases involving zero observed failures for some components where maximum likelihood methods and other proposed approximations tend to break down (see, e.g., Madansky 1965 and Mann et al. 1974). It also permits the pooling of failure data for different components having the same test sample sizes. This potential reduction in the effective number of system components enhances the utility of tabulated bounds such as those presented here.

Because of the structure of the problem, universally optimal confidence bounds for system reliability (i.e., "uniformly most accurate" bounds in the sense of Lehmann 1959) do not generally exist. On the other hand, the ideas of Buehler (1957) may be exploited to produce a variety of valid confidence bounds based on total orderings of the sample points. Such bounds are exact in the sense that the desired coverage probability is guaranteed. The construction of good confidence bounds is thus reduced to the selection of suitable orderings imposed on the sample space. This is the approach adopted in the present study. Previous applications of these methods to reliability may be found, for example, in Harris and Soms (1980), Johnson (1969), and Lipow and Riley (1959).

Once the sample ordering approach has been chosen, there remains the problem of determining orderings which lead to confidence bounds which are "good" according to some criterion measuring the size (length) of the confidence region. An early proposal of the present author (Johns 1975) was to generate the ordering by means of a simple function of the observations which was asymptotically equivalent to the maximum likelihood confidence bound. This method guarantees asymptotic optimality when the numbers of component failures observed under testing is large. It was found, however, that the resulting bounds could be noticeably improved for small to moderate numbers of observed failures by more sophisticated methods. Another procedure investigated (Johns 1977) was a sequential method for generating the sample ordering starting at the origin (zero component failures observed) and selecting at each stage as the next sample

point the "adjacent" point producing the largest value for the lower bound on reliability. This method, while intuitively appealing, does not generally produce a "best" ordering as has been suggested by some investigators. In particular, it is improved upon by the method adopted in the present paper.

The non-existence of a unique optimal ordering leaves open the possibility of obtaining a confidence bound which is at least admissible by choosing the sample ordering to minimize the expected length of the confidence interval under some suitable prior distribution. Such a semi-Bayesian approach does not in any way impair the frequentistic interpretation of the confidence bounds obtained from the minimizing ordering. The implementation of such a minimization, while theoretically perfectly possible, turns out to be totally unfeasible computationally except for the earliest part of the ordering generated. Nevertheless, for a class of priors chosen to emulate certain properties of maximum likelihood, fragmentary orderings computed by this method provide a considerable justification for the two-stage "look-ahead" sequential method actually used to generate the tables which are a principal concern of this paper.

Suppose that the series system under consideration consists of  $k$  independent components and that the respective probabilities of component failure are  $p_i = 1 - q_i$ ,  $i = 1, 2, \dots, k$ . The system reliability  $R$  is given by

$$R = \prod_{i=1}^k q_i = \prod_{i=1}^k (1 - p_i) \quad . \quad (1.1)$$

If the observed numbers of failures for the  $k$  components are  $X_1, X_2, \dots, X_k$  based on independent tests with corresponding sample sizes  $n_1, n_2, \dots, n_k$ , we let  $\lambda_i = n_i p_i$ ,  $i = 1, 2, \dots, k$ , so that

$$R = \prod_{i=1}^k (1 - \lambda_i/n_i) \approx 1 - \sum_{i=1}^k \lambda_i/n_i . \quad (1.2)$$

The approximation on the right will be best when the  $p_i$ 's ( $= \lambda_i/n_i$ ) are all small which is just the case where  $R$  is close to one and the Poisson approximation for the distributions of the  $X_i$ 's is valid.

It will be convenient to express the problem in a canonical form by introducing some further notation. Let  $c = \sum_{i=1}^k 1/n_i$  and  $a_i = 1/cn_i$ ,  $i = 1, 2, \dots, k$ . Then letting  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$  and

$$\theta(\lambda) = \sum_{i=1}^k a_i \lambda_i , \quad (1.3)$$

we have

$$R \approx 1 - c \sum_{i=1}^k a_i \lambda_i = 1 - c\theta(\lambda) . \quad (1.4)$$

We shall assume henceforth that the components are indexed so that  $n_1 \geq n_2 \geq \dots \geq n_k$  which implies that  $a_1 \leq a_2 \leq \dots \leq a_k$ . The problem of finding a lower confidence bound for  $R$  is thus reduced to that of finding an upper confidence bound for  $\theta(\lambda)$ . The fact that  $\theta(\lambda)$  is a convex combination of the  $\lambda_i$ 's facilitates tabulation of the bounds by reducing the number of classification variables by one. In principle, confidence bounds for  $R$  could be constructed directly without introducing the approximation (1.2). Such an approach would,

however, eliminate the possibility of constructing useful tables of bounds, since separate entries would be required for every configuration of  $(n_1, n_2, \dots, n_k)$ .

The vector of observations is  $X = (X_1, X_2, \dots, X_k)$  so that the sample space on which a total ordering must be imposed consists of all vectors  $x = (x_1, x_2, \dots, x_k)$  where the  $x_i$ 's are non-negative integers. It seems reasonable to confine attention to orderings which are consistent with the natural partial ordering induced by dominance (see Section 2), and we shall do so. In Section 2 it is shown that the best upper confidence bound for  $\theta(\lambda)$  with confidence coefficient  $1 - \alpha$  which is monotone in a prescribed total ordering (designated by the relation  $\leq$ ) is given by

$$t(x) = \sup_{\lambda \in S_\alpha^*(x)} \theta(\lambda) , \quad (1.5)$$

where  $S_\alpha^*(x) = \{\lambda : P_\lambda\{X \leq x\} = \alpha\}$ . From (1.4) the lower confidence bound for system reliability is then given by

$$r(x) = 1 - ct(x) . \quad (1.6)$$



Because of the additive property of independent Poisson observations the effective number of components in the system may be reduced if some sample sizes are equal (or nearly equal) as follows: Suppose  $n_1 = n_2 = \dots = n_r = n$  for some  $r$ ,  $2 \leq r < k$ . Then  $a_1 = a_2 = \dots = a_r = a^*$  (say) and letting  $\lambda^* = \sum_{i=1}^r \lambda_i$  we have

$$\theta(\lambda) = a^* \lambda^* + \sum_{i=r+1}^k a_i \lambda_i. \quad (1.7)$$

Now  $X^* = \sum_{i=1}^r X_i$  has a Poisson distribution with parameter  $\lambda^*$  so that the right hand side of (1.7) has the correct form for the case of dimension  $k^* = k - r + 1$  except that the coefficients must each be divided by  $c_0 = a^* + a_{r+1} + \dots + a_k$  to preserve convexity. The confidence bound for the  $k^*$  dimensional case based on  $X^*, X_{r+1}, \dots, X_k$  and the normalized coefficients may then be multiplied by  $c_0$  to obtain the bound for the original  $\theta(\lambda)$  given by (1.7). Further reductions may be made in the same way if several groups of components have common sample sizes. If  $n_1, n_2, \dots, n_r$  are only approximately equal, their average may be used for  $n$  in the above calculations to obtain an approximate bound.

If all components are subjected to the same number of trials, we may take  $k$  equal to one and the problem reduces to the familiar one of finding an upper confidence bound for a single Poisson parameter.

The component failure data may be developed through independent testing of the components, or through testing of the complete system with the assignment of failures to the appropriate components.

Even in the latter case component sample sizes may differ if components are redesigned during the course of testing so that the trials and failures observed prior to redesign are not relevant to the reliability of the final version of the system.

It is worth noting that the same formal confidence bound problem arises from a somewhat different and more restrictive model involving continuous time sampling and exponentially distributed failure times. Thus, if we assume that the time to failure of the  $i$ -th component has the exponential probability density function  $\mu_i e^{-\mu_i t}$ ,  $\mu_i > 0$ ,  $t > 0$ , and if the fixed aggregate testing time for the  $i$ -th component is given by  $\tau_i$  and the total number of failures observed is  $X_i$ ,  $i=1,2,\dots,k$ , then the  $X_i$ 's which are sufficient statistics have independent Poisson distributions with corresponding parameters  $\lambda_i = \mu_i \tau_i$ . Such test data would be generated, for example, if a single unit of each component were placed on test and repaired whenever a failure occurred until the total test time for each component reached its corresponding limit,  $\tau_i$ . The system reliability for a single duty cycle (consisting of operation for a unit interval of time) is given by

$$R = \prod_{i=1}^k e^{-\mu_i} = \exp \left\{ - \sum_{i=1}^k \mu_i \right\} = \exp \left\{ - \sum_{i=1}^k \lambda_i / \tau_i \right\}, \quad (1.8)$$

and the confidence bound problem is essentially the same as the one previously introduced except that no approximations are required.

The general theory of confidence bounds based on sample orderings is discussed in Section 2. In Section 3 the case of systems having two components is considered in detail, and the rationale for, and use of, the tables for this case are explained. Section 4 includes suggestions for constructing approximate confidence bounds for cases of systems with  $k \geq 3$  by finding approximately equivalent cases with  $k = 2$ . Use of the maximum likelihood ratio bounds for cases where the data are beyond the limits of the available tables is also discussed.

## 2. Bounds and Orderings: Generalities

The idea of using sample orderings to generate confidence bounds was first introduced by Buehler (1957) who discussed the validity of the proposed method in the context of a specific reliability problem. Bol'shev and Loginov (1969) discuss the construction of confidence bounds monotone in the sample orderings generated by certain functions of the observations. In the following, which is a revision and extension of Johns (1975), we develop the theory with emphasis on the sample orderings themselves rather than possible generators of the orderings.

To develop the general ideas relating exact confidence bounds to sample orderings it is convenient to introduce a fairly abstract statistical model. Let the sample space  $\mathcal{X}$  be endowed with a measurable total ordering relation " $\leq$ " and let  $X$  represent the random outcome of the experiment where the space of possible outcomes is  $\mathcal{X}$ . Suppose that the possible distributions of  $X$  are determined by the family of probability measures  $P_\lambda$ , indexed by  $\lambda$ , an element of the parameter space  $\Lambda$ . Our objective is to find a  $1 - \alpha$  level upper confidence bound for a specified real-valued function  $\theta(\lambda)$  defined on  $\Lambda$  where the range  $T$  of  $\theta(\lambda)$  is assumed to be closed and bounded below. The quantity  $\alpha \in (0,1)$  is regarded as fixed throughout. We make the following definitions and assumptions:

Definition D1. For each  $x \in \mathcal{X}$  let

$$S_\alpha(x) = \{\lambda : P_\lambda\{X \leq x\} > \alpha\} . \quad (2.1)$$

Definition D2. For each  $x \in \mathcal{X}$  let

$$t(x) = \begin{cases} \sup\{\theta(\lambda) : \lambda \in S_\alpha(x)\} , & \text{if } S_\alpha(x) \text{ is non-empty} , \\ \inf T, & \text{otherwise} . \end{cases} \quad (2.2)$$

Remark 1. By D2 if  $\theta(\lambda) > t(x)$ , then necessarily  $P_\lambda\{X \leq x\} \leq \alpha$ .

Assumption A1. For every subset  $C$  of  $\mathcal{X}$  having the property that if  $x \in C$  and  $y \leq x$  then  $y \in C$ , there exists an ordered sequence

$x_1 \leq x_2 \leq \dots$  of elements of  $C$  such that  $C = \bigcup_{n=1}^{\infty} \{x : x \leq x_n\}$ .

Assumption A2. For each  $x \in \mathcal{X}$ , if  $\theta(\lambda) = t(x)$ , then  $P_\lambda\{X \leq x\} \leq \alpha$ .

Remark 2. By D1, D2, and A2 if  $\theta(\lambda) = t(x)$ , then  $\lambda \notin S_\alpha(x)$ , i.e., the supremum in D2 is never attained.

We now establish the following propositions.

Proposition P1. The function  $t(x)$  is monotone in the ordering on  $\mathcal{X}$ .

Proof: If  $x, y \in \mathcal{X}$  and  $x \leq y$ , then  $S_\alpha(x) \subset S_\alpha(y)$  (D1) which in turn implies  $t(x) \leq t(y)$  (D2).  $\square$

Proposition P2. Under assumptions A1 and A2 the function  $t(x)$  is an upper confidence bound for  $\theta(\lambda)$  at level  $1 - \alpha$ . In particular,

$$P_\lambda \{ \theta(\lambda) < t(X) \} \leq \alpha \quad \text{for all } \lambda \in \Lambda. \quad (2.3)$$

Proof: For arbitrary  $\lambda_0 \in \Lambda$ , let  $\theta_0 = \theta(\lambda_0)$  and  $C_0 = \{x: t(x) \leq \theta_0\}$ .

The result follows immediately for all  $\lambda_0$  for which  $C_0$  is empty.

Assume that  $C_0$  is non-empty. Then by P1 the set  $C_0$  possesses the property required in A1 for the existence of a sequence  $\{x_n\} \subset C_0$  such that  $x_n \leq x_{n+1}$  for all  $n$ , and  $C_0 = \bigcup_{n=1}^{\infty} \{x: x \leq x_n\}$ . This implies that, as  $n \rightarrow \infty$ ,

$$P_{\lambda_0} \{X \leq x_n\} \uparrow P_{\lambda_0} \{X \in C_0\}. \quad (2.4)$$

But by Remark 1 and A2, for all  $n$ ,  $P_{\lambda_0} \{X \leq x_n\} < \alpha$ . Hence

$P_{\lambda_0} \{X \in C_0\} \leq \alpha$  and the desired result follows.  $\square$

Proposition P3. Under assumptions A1 and A2, if  $\tilde{t}(x)$  is any T-valued confidence bound such that  $P_\lambda\{\theta(\lambda) < \tilde{t}(X)\} \geq 1 - \alpha$  for all  $\lambda \in \Lambda$ , then

$$(i) \quad \sup_{y \leq x} \tilde{t}(y) \geq t(x) \quad \text{for all } x \in \mathcal{X}, \quad \text{and}$$

(ii) if  $t(x)$  is monotone in the ordering on  $\mathcal{X}$ , then

$$\tilde{t}(x) \geq t(x) \quad \text{for all } x \in \mathcal{X}.$$

Proof: First we assume that  $\tilde{t}(x)$  is monotone and establish (ii).

Suppose there exists an  $x' \in \mathcal{X}$  such that  $\tilde{t}(x') < t(x')$ . Then  $S_\alpha(x')$  must be non-empty and by Remark 2 following A2, the sup defining  $t(x')$  is not attained. Hence there exists a  $\lambda' \in S_\alpha(x')$  such that  $\tilde{t}(x') < \theta(\lambda') < t(x')$ , and  $P_{\lambda'}\{X \leq x'\} > \alpha$ . Thus by the monotonicity of  $\tilde{t}(x)$ ,

$$P_{\lambda'}\{\tilde{t}(X) \leq \theta(\lambda')\} \geq P_{\lambda'}\{\tilde{t}(X) \leq \tilde{t}(x')\} = P_{\lambda'}\{X \leq x'\} > \alpha. \quad (2.5)$$

This contradicts the hypothesis that  $P_\lambda\{\theta(\lambda) < \tilde{t}(X)\} \geq 1 - \alpha$  for all  $\lambda$  and establishes (ii). To show (i) we let  $t^*(x) = \sup_{y \leq x} \tilde{t}(y)$  and observe that  $t^*(x)$  is monotone in the ordering on  $\mathcal{X}$  and  $t^*(x) \geq \tilde{t}(x)$  for all  $x \in \mathcal{X}$ . Hence if  $\tilde{t}(x)$  is a  $1 - \alpha$  confidence bound for  $\theta(\lambda)$ , so is  $t^*(x)$  and applying (ii) to  $t^*(x)$  yields (i).  $\square$ .

In order to specialize these results in the direction of applications we henceforth assume that the parameter  $\lambda$  and the

observation  $X$  are both of dimension  $k$ , i.e.,  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$  and  $X = (X_1, X_2, \dots, X_k)$  where the  $\lambda_i$ 's are real and the  $X_i$ 's are random variables. Without essential loss of generality we assume that  $\Lambda$  contains the positive orthant. Within this framework we make the following additional assumptions:

Assumption A3. The function  $\theta(\lambda)$  is continuous and strictly increasing in each of the  $\lambda_i$ 's.

Assumption A4. For any  $x \in \mathcal{X}$ ,  $P_\lambda\{X \leq x\}$  is continuous in each of the  $\lambda_i$ 's.

Proposition P4. Assumptions A3 and A4 imply that Assumption A2 is satisfied.

Proof: Suppose that for some  $x' \in \mathcal{X}$  there exists a  $\lambda' \in \Lambda$  such that  $\theta(\lambda') = t(x')$  and  $P_{\lambda'}\{X \leq x'\} > \alpha$ . Then by A3 and A4 we can find a  $\lambda'' \in \Lambda$  with  $\lambda_i'' \geq \lambda_i'$  for all  $i$  and  $\lambda_{i_0}'' > \lambda_{i_0}'$  for some  $i_0$  such that  $\theta(\lambda'') > t(x')$  and  $P_{\lambda''}\{X \leq x'\} > \alpha$ . This contradicts D2 and the result follows.  $\square$

Corollary C1. If (i) the observations  $X_1, X_2, \dots, X_k$  are independent Poisson random variables with parameters  $\lambda_1, \lambda_2, \dots, \lambda_k$  respectively, and (ii)  $\theta(\lambda) = a_1\lambda_1 + a_2\lambda_2 + \dots + a_k\lambda_k$  where the  $a_i$ 's are positive, then  $t(x)$  given by D2 is an upper confidence bound for  $\theta(\lambda)$  at level  $1 - \alpha$ .

Proof: A1 is satisfied for any ordering since  $\mathcal{X}$  is discrete. A3 is clearly satisfied for  $\theta(\lambda)$  of the form given by (ii), and A4 is satisfied for Poisson random variables. The desired result follows from P4 and P2.  $\square$

The actual computation of the bound  $t(x)$  given by D2 is greatly facilitated if  $P_\lambda\{X \leq x\}$  is monotone in the components of  $\lambda$ . The following proposition gives conditions guaranteeing this property:

Assumption A5. The components of  $X$  are independent and for  $i=1,2,\dots,k$  the distribution of each  $X_i$  depends only on the corresponding  $\lambda_i$  and is stochastically increasing in  $\lambda_i$ .

Definition D3. We denote by  $(\preceq^*)$  the natural partial ordering of  $\mathcal{X}$  generated by componentwise dominance. That is,  $x \preceq^* y$  if and only if  $x_i \leq y_i$  for  $i=1,2,\dots,k$ , with  $x \prec^* y$  if at least one of these inequalities is strict. An arbitrary total ordering  $(\preceq)$  on  $\mathcal{X}$  is consistent with  $(\preceq^*)$  if  $x \preceq^* y$  implies  $x \preceq y$ .

Proposition P5. If the ordering on  $\mathcal{X}$  is consistent with the natural partial ordering and Assumption A5 is satisfied, then for each  $x \in \mathcal{X}$ ,  $P_\lambda\{X \leq x\}$  is non-increasing in each component of  $\lambda$ .

Proof: For  $y \in \mathcal{X}$  we introduce the representation  $y = (y_1, y^{(2)})$  where  $y^{(2)} = (y_2, y_3, \dots, y_k)$ . For real  $z$ , fixed  $x \in \mathcal{X}$ , and all  $y \in \mathcal{X}$ , let  $I_{x,z,y^{(2)}}$  be the indicator function of the set  $\{y^{(2)} : (z, y^{(2)}) \leq x\}$ .



Then

$$P_{\lambda}\{X \leq x\} = E_{\lambda} I_x(X_1, X^{(2)}) \quad (2.6)$$

For any  $y \in \mathcal{X}$ , if  $z' \leq z''$ , then  $(z', y^{(2)}) \leq^* (z'', y^{(2)})$ , by the consistency hypothesis, and  $I_x(z', y^{(2)}) \geq I_x(z'', y^{(2)})$ . Hence letting  $G_x(z) = E_{\lambda}\{I_x(X_1, X^{(2)}) | X_1 = z\}$  we see that  $G_x(z)$  is non-increasing in  $z$ . Thus, since the distribution of  $X_1$  is stochastically increasing in  $\lambda_1$  (A5), we conclude that  $E_{\lambda} I_x(X_1, X^{(2)}) = E_{\lambda} G_x(X_1)$  is non-increasing in  $\lambda_1$ . The same argument applies to the other components of  $\lambda$  establishing the desired result.  $\square$

Suppose that  $\Lambda$  is the non-negative orthant of  $R^{(k)}$  and let  $\mathcal{S}$  be the simplex  $\mathcal{S} = \{\lambda : \sum_{i=1}^k \lambda_i = 1\}$ . If A4 and A5 are satisfied, then by P5 we observe that for any  $\lambda \in \mathcal{S}$  and real  $c$ ,  $P_{c\lambda}\{X \leq x\}$  is continuous and non-increasing in  $c$ . If the lower bound of  $P_{c\lambda}\{X \leq x\}$  as  $c \rightarrow \infty$  is less than  $\alpha$  for all  $x \in \mathcal{X}$  and all  $\lambda \in \mathcal{S}$ , then for each  $x \in \mathcal{X}$  and  $\lambda \in \mathcal{S}$  there exists a smallest number  $b = b(x, \lambda)$  such that  $P_{b(x, \lambda)\lambda}\{X \leq x\} = \alpha$ . The confidence bound  $t(x)$  defined by D2 is then given by

$$t(x) = \sup_{\lambda \in \mathcal{S}} \theta(b(x, \lambda)\lambda) \quad (2.7)$$

Now  $b(x, \lambda)$  is easily computed using root-finding techniques so that the computation of  $t(x)$  reduces to searching over  $\mathcal{S}$  for the maximum of  $\theta(b(x, \lambda)\lambda)$ . Many routines are available for implementing such searches. For the situation described in Corollary C1, the value of

b such that  $P_{b\lambda}\{X \leq x\} = \alpha$  is unique and (2.7) is a computationally feasible version of (1.5).

All of the above results apply mutatis mutandis to the construction of lower confidence bounds and hence confidence intervals. Applications to the reliability of coherent systems involving the binomial or other distributions are possible. In particular, the above discussion applies directly to the binomial case under the transformation  $\lambda_i = -\log(1-p_i)$ ,  $i = 1, 2, \dots, k$ ;  
 $\theta(\lambda) = \sum_{i=1}^k \lambda_i = -\log \prod_{i=1}^k (1-p_i)$ .

### 3. Systems With $k = 2$

As was noted in Section 1, if the system has effectively only one component (e.g., when all sample sizes are equal), then the problem reduces to the well-known case of finding an upper confidence bound for a single Poisson parameter. Then in the notation of Section 1,  $\theta(\lambda) = \lambda$  and if  $t(x)$  is the confidence bound for  $\lambda$ , the lower confidence bound (1.6) for reliability  $R$  becomes

$$r(x) = 1 - t(x)/n, \quad (3.1)$$

where  $n$  is the (common) sample size.

The two component case ( $k = 2$ ) presents all of the difficulties of the general case. The principal problem is to generate an ordering of the sample points  $x = (x_1, x_2)$  which will lead to a "good" confidence bound  $t(x)$  computed using (1.5) or (2.7). Several different methods have been considered and implemented to varying extents

in the course of this investigation. These methods may be described briefly as follows:

- (i) The  $x$ 's are ordered according to the values of the function  $\tilde{t}(x) = a_1x_1 + a_2x_2 + z_\alpha \sqrt{a_1^2x_1^2 + a_2^2x_2^2}$ , where  $z_\alpha$  is the upper  $\alpha$ -th quantile of the standard normal distribution.
- (ii) The  $x$ 's are ordered according to the values of the approximate confidence bound obtained from the maximum likelihood ratio statistic (see Section 4).
- (iii) The ordering is generated sequentially by considering at each stage the group of points which are not yet ordered but could be adjoined without violating the natural partial ordering (see D3 of Section 2). The next point in the ordering is then chosen to be the "best" member of the candidate group, i.e., the point producing the smallest value of  $t(x)$  given by (1.5).
- (iv) The ordering is chosen so as to minimize  $E_G\{t(X)\}$  for some suitable prior distribution  $G$  over the values of  $\lambda$ .
- (v) The ordering is generated sequentially in the manner of (iii) above except that at each stage the candidate points for the next two steps are considered as pairs and the next point selected is the first step component which, together with the best available point for the second step, produces the smallest sum for the two values of  $t(x)$ . Note that the point that appears to be "best" two steps ahead may not actually be chosen when that stage is reached.

It is clear that none of these methods is special to the case  $k = 2$ . Method (i), based on the function  $\tilde{t}(x)$ , which is really a maximum likelihood estimate of an asymptotically valid confidence bound, was used to generate tables of bounds for the case  $k = 2$  in Johns (1975). Method (ii) does not improve substantially on Method (i) for moderate values of the  $X_i$ 's. Methods (i) and (ii) are asymptotically equivalent when at least one  $X_i$  becomes large (see Johns 1975) and indeed standard maximum likelihood results guarantee that both are asymptotically optimal. Method (iii) discussed in Johns (1977) was found to be a substantial improvement on (i) in the strong sense that when the Method (iii) ordering is used the values of  $t(x)$  are often smaller and only rarely slightly larger than the values for corresponding  $x$ 's produced by Method (i).

The semi-Bayesian approach of Method (iv), which minimizes the expected length of the confidence interval, is the only one of the five that is directly motivated by optimality considerations. The bound resulting from any reasonable prior must at least be admissible. In pursuing this approach it was decided in the spirit of objectivity and in the hope of rapid convergence to asymptotic optimality to choose a prior distribution leading to an unconditional probability mass function for  $x_1$  and  $x_2$  constant for constant values of the maximum likelihood estimator  $a_1x_1 + a_2x_2$  for  $\theta(\lambda)$ . In particular, the prior density for  $\lambda_1$  and  $\lambda_2$  was taken to be

$$g(\lambda_1, \lambda_2) = b_1 b_2 e^{-b_1 \lambda_1 - b_2 \lambda_2}, \quad \lambda_1, \lambda_2 > 0, \quad (3.2)$$

where  $b_1 = (1 - e^{-\beta a_1})$ ,  $b_2 = (1 - e^{-\beta a_2})$ ,  $\beta > 0$ . This produces the unconditional probability mass function

$$p(x_1, x_2) = b_1 b_2 e^{-\beta(a_1 x_1 + a_2 x_2)}, \quad x_1, x_2 = 0, 1, \dots, \quad (3.3)$$

In the limiting case, as  $\beta \rightarrow 0$ ,  $p(x_1, x_2)$  becomes essentially uniform over any finite set of points  $(x_1, x_2)$ .

The actual minimization of  $E_G\{t(X)\}$  may, in principle, be accomplished by finding the ordering which minimizes the contribution to  $E_G$  among all orderings of length  $N$  where  $N$  may be arbitrarily large. This may be done systematically by starting at the origin  $(0,0)$  and constructing a tree whose nodes at each stage are characterized by a candidate point newly adjoined to the ordering and the corresponding value of  $\sum p(x)t(x)$ , where the sum is taken over all  $x$ 's occurring in the path leading to the node, including the one just adjoined. At the  $N$ -th stage the node having the smallest accumulated sum identifies the optimal ordering of length  $N$ . This process may be facilitated by eliminating duplicate nodes and discontinuing branches when a node is reached whose value exceeds that known to be attainable in  $N$  stages. Nevertheless, because of the rapid increase in the number of nodes considered per stage, only the first forty or so points in the optimal orderings could be determined even using a very large computer facility.

In order to obtain examples of admissible orderings with which to compare the results of other methods, this computation was performed for two cases using an IBM 370/168. For both cases the

values  $\alpha = .10$  and  $a_1 = .30$  were used. For the first case the probabilities given in (3.3) with  $\beta = 1$  were used and the first 41 points of the optimal ordering were obtained. The forty-first stage of the computation produced 4557 nodes. For the second case the limiting situation as  $\beta \rightarrow 0$  where the  $p(x)$ 's are all equal was used and the first 43 points of the optimal ordering were obtained. The number of nodes produced at the forty-third stage was 6478.

A comparison of these results with the corresponding results obtained using Methods (i), (iii), and (v) is indicated in Figure 1. The horizontal axis indexes the first 50 points in the ordering produced by Method (v), the two-step prospective sequential procedure. The values of  $t(x)$  for these indexed points for the five methods are indicated by the plotted symbols. Values of  $t(x)$  for methods other than (v) are shown only when they differ from those produced by that method. Based on this evidence it appears that Method (iv) and Method (v) differ very little and that both are better than the other methods. In fact, Method (iv) for the uniform case ( $\beta = 0$ ) differs only trivially from Method (v). Since the use of Method (iv) for the construction of tables is now and probably always will be impractical, we are led to the choice for this purpose of the more tractable and virtually equivalent Method (v). Prospective sequential methods looking ahead more than two steps might be feasible, although the complexity of the computations increases rapidly with the number of steps. However, such procedures would be expected to produce only minute improvement over the two-step method.




Figure 1. Comparison of confidence bounds for five ordering methods:  $\alpha = .10$ ,  $a_1 = .30$

Table 1 gives values of the upper confidence bound  $t(x)$  computed using (2.7) with the sample orderings generated by the two-step prospective sequential Method (v). The values are given for the first 100 points in each ordering for  $\alpha = .01, .05, .10$ , for  $a_1 = .05(.05).45$ . It is assumed that the components of the system are indexed so that  $a_1 < a_2$  which implies  $a_1 < .50$ . For convenience the values of  $(x_1, x_2)$  are listed systematically rather than in the order generated by the two-step procedure. This table provides a basis for computing accurate confidence bounds for the case  $k = 2$  using only simple interpolation. If values of  $a_1$  greater than .45 but (necessarily) less than .50 are required, the bound for  $a_1 = .5$  (corresponding to  $n_1 = n_2$ ) may be used for interpolation. This bound is obtained by simply multiplying the ordinary upper confidence bound for a single Poisson parameter based on  $x = x_1 + x_2$  failures by .50 (see, e.g., Pearson and Hartley 1958 for tables). The use of Table 1 is illustrated by the following two examples:

Example 1. Suppose that the two components of a series system are tested independently using sample sizes  $n_1 = 300$  and  $n_2 = 100$  respectively with the corresponding observed numbers of failures  $X_1 = 3$  and  $X_2 = 4$ . Then  $c = (1/300 + 1/100) = 4/300$  and  $a_1 = 1/cn_1 = .25 = 1 - a_2$ . If we wish to find a 95 percent confidence interval, we take  $\alpha = .05$ , and from Table 1 we find the confidence bound  $t(x)$  for  $\theta(\lambda)$  to be 7.333. Hence by (1.6) the 95 percent lower confidence bound for system reliability  $R$  is  $1 - (4/300)(7.333) = .902$ .



Table 1. The Confidence Bound  $t(x)$  for  $k = 2$  for the First 100 Points Generated by the Two-Stage Optimal Ordering Method for Each  $a_1$  and  $\alpha$ .

$a_1 = 0.05$				$a_1 = 0.05$				$a_1 = 0.05$				$a_1 = 0.05$				$a_1 = 0.10$			
$x_1$	$x_2$	$t(x_1, x_2)$	$\alpha$	$x_1$	$x_2$	$t(x_1, x_2)$	$\alpha$	$x_1$	$x_2$	$t(x_1, x_2)$	$\alpha$	$x_1$	$x_2$	$t(x_1, x_2)$	$\alpha$	$x_1$	$x_2$	$t(x_1, x_2)$	$\alpha$
0	0	4.375	25	0	0	2.846	25	0	0	3.733	25	0	0	2.188	25	0	0	3.075	25
0	1	6.306	25	0	1	4.507	25	0	1	5.396	25	0	1	3.695	25	0	1	4.585	25
1	0	4.376	26	1	0	2.847	26	1	0	3.777	26	1	0	2.189	26	1	0	3.119	26
1	1	6.308	26	1	1	4.508	26	1	1	5.440	26	1	1	3.697	26	1	1	4.629	26
2	0	4.387	27	2	0	2.858	27	2	0	3.822	27	2	0	2.200	27	2	0	3.163	27
2	1	6.319	27	2	1	4.519	27	2	1	5.485	27	2	1	3.708	27	2	1	4.674	27
3	0	4.407	28	3	0	2.878	28	3	0	3.866	28	3	0	2.220	28	3	0	3.208	28
3	1	6.327	28	3	1	4.539	28	3	1	5.529	28	3	1	3.727	28	3	1	4.718	28
4	0	4.432	29	4	0	2.903	29	4	0	3.911	29	4	0	2.245	29	4	0	3.253	29
4	1	6.364	29	4	1	4.564	29	4	1	5.574	29	4	1	3.753	29	4	1	4.763	29
5	0	4.461	30	5	0	2.932	30	5	0	3.956	30	5	0	2.274	30	5	0	3.298	30
5	1	6.393	30	5	1	4.593	30	5	1	5.619	30	5	1	3.781	30	5	1	4.809	30
6	0	4.493	31	6	0	2.964	31	6	0	4.001	31	6	0	2.305	31	6	0	3.343	31
6	1	6.424	31	6	1	4.624	31	6	1	5.665	31	6	1	3.813	31	6	1	4.854	31
7	0	4.526	32	7	0	2.997	32	7	0	4.047	32	7	0	2.339	32	7	0	3.388	32
7	1	6.458	32	7	1	4.658	32	7	1	5.692	32	7	1	3.847	32	7	1	4.899	32
8	0	4.561	33	8	0	3.032	33	8	0	4.138	33	8	0	2.374	33	8	0	3.434	33
8	1	6.493	33	8	1	4.693	33	8	1	5.735	33	8	1	3.882	33	8	1	4.945	33
9	0	4.598	34	9	0	3.069	34	9	0	4.229	34	9	0	2.410	34	9	0	3.479	34
9	1	6.529	34	9	1	4.730	34	9	1	5.771	34	9	1	3.918	34	9	1	4.985	34
10	0	4.635	35	10	0	3.106	35	10	0	4.321	35	10	0	2.447	35	10	0	3.525	35
10	1	6.567	35	10	1	4.767	35	10	1	5.803	35	10	1	3.956	35	10	1	5.011	35
11	0	4.673	36	11	0	3.144	36	11	0	4.413	36	11	0	2.486	36	11	0	3.571	36
11	1	6.605	36	11	1	4.805	36	11	1	5.835	36	11	1	3.994	36	11	1	5.039	36
12	0	4.712	37	12	0	3.183	37	12	0	4.506	37	12	0	2.525	37	12	0	3.616	37
12	1	6.644	37	12	1	4.845	37	12	1	5.867	37	12	1	4.033	37	12	1	5.071	37
13	0	4.752	38	13	0	3.223	38	13	0	4.599	38	13	0	2.565	38	13	0	3.647	38
13	1	6.684	38	13	1	4.884	38	13	1	5.899	38	13	1	4.073	38	13	1	5.101	38
14	0	4.792	39	14	0	3.264	39	14	0	4.692	39	14	0	2.605	39	14	0	3.684	39
14	1	6.724	39	14	1	4.925	39	14	1	5.929	39	14	1	4.114	39	14	1	5.131	39
15	0	4.833	40	15	0	3.304	40	15	0	4.739	40	15	0	2.646	40	15	0	3.721	40
15	1	6.765	40	15	1	4.966	40	15	1	5.961	40	15	1	4.154	40	15	1	5.161	40
16	0	4.875	41	16	0	3.346	41	16	0	4.833	41	16	0	2.687	41	16	0	3.755	41
16	1	6.807	41	16	1	5.007	41	16	1	6.003	41	16	1	4.196	41	16	1	5.191	41
17	0	4.916	42	17	0	3.388	42	17	0	4.974	42	17	0	2.729	42	17	0	3.789	42
17	1	6.849	42	17	1	5.049	42	17	1	6.035	42	17	1	4.238	42	17	1	5.221	42
18	0	4.959	43	18	0	3.430	43	18	0	5.068	43	18	0	2.771	43	18	0	3.827	43
18	1	6.891	43	18	1	5.091	43	18	1	6.068	43	18	1	4.280	43	18	1	5.251	43
19	0	5.001	44	19	0	3.472	44	19	0	5.163	44	19	0	2.814	44	19	0	3.864	44
19	1	6.933	44	19	1	5.134	44	19	1	6.100	44	19	1	4.323	44	19	1	5.281	44
20	0	5.044	45	20	0	3.515	45	20	0	5.210	45	20	0	2.856	45	20	0	3.901	45
20	1	6.976	45	20	1	5.177	45	20	1	6.135	45	20	1	4.366	45	20	1	5.311	45
21	0	5.087	46	21	0	3.558	46	21	0	5.305	46	21	0	2.900	46	21	0	3.940	46
21	1	7.019	46	21	1	5.220	46	21	1	6.160	46	21	1	4.409	46	21	1	5.341	46
22	0	5.130	47	22	0	3.601	47	22	0	5.400	47	22	0	2.943	47	22	0	3.979	47
22	1	7.063	47	22	1	5.264	47	22	1	6.191	47	22	1	4.453	47	22	1	5.381	47
23	0	5.174	48	23	0	3.645	48	23	0	5.495	48	23	0	2.987	48	23	0	4.019	48
23	1	7.107	48	23	1	5.307	48	23	1	6.232	48	23	1	4.496	48	23	1	5.421	48
24	0	5.218	49	24	0	3.689	49	24	0	5.591	49	24	0	3.031	49	24	0	4.059	49
24	1	7.151	49	24	1	5.351	49	24	1	6.273	49	24	1	4.540	49	24	1	5.461	49

Table 1. (Continued)

a1 = 0.1    alpha = 0.01				a1 = 0.1    alpha = 0.05				a1 = 0.1    alpha = 0.10						
x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)
0	0	4.145	17	0	5.384	0	0	2.696	16	1	5.426	0	0	2.072
0	1	5.975	17	1	7.221	0	1	4.269	16	2	6.827	0	1	3.501
0	2	7.565	18	0	5.477	0	2	5.666	17	0	3.936	0	2	4.790
1	0	4.151	18	1	7.314	0	3	6.978	17	1	5.519	0	3	6.013
1	1	5.981	19	0	5.569	1	0	2.702	17	2	6.920	0	4	4.659
1	2	7.571	19	1	7.407	1	1	4.275	18	0	4.028	1	1	3.507
2	0	4.184	20	0	5.662	1	2	5.672	18	1	5.612	1	2	4.796
2	1	6.014	20	1	7.501	2	0	2.735	19	0	4.121	1	3	6.019
2	2	7.605	21	0	5.756	2	1	4.309	19	1	5.706	2	0	2.111
3	0	4.235	21	1	7.596	2	2	5.706	20	0	4.214	2	1	3.540
3	1	6.065	22	0	5.850	3	0	2.787	20	1	5.800	2	2	4.829
3	2	7.656	22	1	7.691	3	1	4.360	21	0	4.307	2	3	6.052
4	0	4.297	23	0	5.945	3	2	5.757	21	1	5.895	3	0	2.163
4	1	6.127	23	1	7.786	4	0	2.849	22	0	4.402	3	1	3.590
4	2	7.718	24	0	6.040	4	1	4.422	22	1	5.991	2	2	4.881
5	0	4.366	24	1	7.882	4	2	5.819	23	0	4.496	3	3	6.104
5	1	6.196	25	0	6.135	5	0	2.917	23	1	6.087	4	0	2.225
5	2	7.786	25	1	7.978	5	1	4.491	24	0	4.591	4	1	3.654
6	0	4.439	26	0	6.230	5	2	5.889	24	1	6.183	4	2	4.943
6	1	6.270	26	1	8.075	6	0	2.991	25	0	4.687	5	0	2.293
6	2	7.861	27	0	6.326	6	1	4.565	25	1	6.280	5	1	3.723
7	0	4.516	27	1	8.172	6	2	5.963	26	0	4.782	5	2	5.013
7	1	6.348	28	0	6.422	7	0	3.068	26	1	6.380	6	0	2.367
7	2	7.939	28	1	8.270	7	1	4.643	27	0	4.878	6	1	3.797
8	0	4.596	29	0	6.519	7	2	6.040	27	1	6.478	6	2	5.087
8	1	6.428	29	1	8.369	8	0	3.148	28	0	4.975	7	0	2.444
8	2	8.020	30	0	6.616	8	1	4.723	28	1	6.577	7	1	3.875
9	0	4.679	30	1	8.467	8	2	6.121	29	0	5.073	7	2	5.165
9	1	6.510	31	0	6.713	9	0	3.230	29	1	6.676	8	0	2.524
9	2	8.102	31	1	8.565	9	1	4.806	30	0	5.170	8	1	3.955
10	0	4.763	32	0	6.810	9	2	6.204	30	1	6.776	8	2	5.246
10	1	6.595	32	1	8.665	10	0	3.314	31	0	5.268	9	0	2.606
10	2	8.187	33	0	6.908	10	1	4.891	31	1	6.878	9	1	4.038
1	0	4.681	34	0	7.006	10	2	6.289	32	0	5.366	9	2	5.329
1	1	6.681	35	0	7.104	11	0	3.400	33	0	5.464	10	0	2.690
1	2	8.273	36	0	7.202	11	1	4.977	34	0	5.563	10	1	4.123
2	0	4.935	37	0	7.300	11	2	6.376	35	0	5.661	10	2	5.414
2	1	6.768	38	0	7.399	12	0	3.487	36	0	5.760	11	0	2.776
2	2	8.361	39	0	7.498	12	1	5.065	37	0	5.859	11	1	4.209
3	0	5.023	40	0	7.600	12	2	6.464	38	0	5.959	11	2	5.501
3	1	6.857	41	0	7.700	13	0	3.575	39	0	6.058	12	0	2.863
3	2	8.450	42	0	7.799	13	1	5.154	40	0	6.158	12	1	4.297
4	0	5.112	43	0	7.899	13	2	6.553	41	0	6.258	12	2	5.590
4	1	6.947	44	0	8.000	14	0	3.664	42	0	6.359	13	0	2.951
4	2	8.540	45	0	8.100	14	1	5.244	43	0	6.459	13	1	4.386
5	0	5.202	46	0	8.200	14	2	6.644	44	0	6.559	13	2	5.679
5	1	7.037	47	0	8.301	15	0	3.754	45	0	6.660	14	0	3.040
5	2	8.631	48	0	8.402	15	1	5.334	46	0	6.761	14	1	4.476
6	0	5.293	49	0	8.503	15	2	6.735	47	0	6.862	14	2	5.770
6	1	7.129	50	0	8.604	16	0	3.844	48	0	6.963	15	0	3.130
6	2													
7	0													
7	1													
7	2													
8	0													
8	1													
8	2													
9	0													
9	1													
9	2													
10	0													
10	1													
10	2													
1	0													
1	1													
1	2													
2	0													
2	1													
2	2													
3	0													
3	1													
3	2													
4	0													
4	1													
4	2													
5	0													
5	1													
5	2													
6	0													
6	1													
6	2													
7	0													
7	1													
7	2													
8	0													
8	1													
8	2													
9	0													
9	1													
9	2													
10	0													
10	1													
10	2													
1	0													
1	1													
1	2													
2	0													
2	1													
2	2													
3	0													
3	1													
3	2													
4	0													
4	1													
4	2													
5	0													
5	1													
5	2													
6	0													
6	1													
6	2													
7	0													
7	1													
7	2													
8	0													
8	1													
8	2													
9	0													
9	1													
9	2													
10	0													
10	1													
10	2													
1	0													
1	1													
1	2													
2	0													
2	1													
2	2													
3	0													
3	1													
3	2													
4	0													
4	1													
4	2													
5	0													
5	1													
5	2													
6	0													
6	1													
6	2													
7	0													
7	1													
7	2													

Table 1. (Continued)

a1 = 0.15				alpha = 0.01				a1 = 0.15				alpha = 0.05				a1 = 0.15				alpha = 0.10			
x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)
0	0	3.914	7.143	13	1	7.143	0	0	2.546	12	0	3.883	0	0	1.957	11	2	5.755	11	2	5.755		
0	1	5.643	8.659	13	2	8.659	12	1	4.032	12	1	5.403	0	1	3.306	11	3	6.921	11	3	6.921		
0	2	7.145	5.538	14	0	5.538	12	2	5.351	12	2	6.728	0	2	4.524	12	0	3.293	12	0	3.293		
0	3	8.538	7.291	14	1	7.291	13	0	6.591	13	0	4.026	0	3	5.679	12	1	4.679	12	1	4.679		
1	0	3.929	8.809	14	2	8.809	13	1	7.780	13	1	5.560	0	4	6.795	12	2	5.905	12	2	5.905		
1	1	5.658	5.684	15	0	5.684	13	2	2.561	13	2	5.681	0	0	1.972	13	0	3.438	13	0	3.438		
1	2	7.160	7.443	15	1	7.443	14	0	4.047	14	0	4.171	1	1	3.321	13	1	4.836	13	1	4.836		
1	3	8.553	8.964	15	2	8.964	14	1	5.367	14	1	5.717	1	2	4.530	13	2	6.055	13	2	6.055		
2	0	3.994	5.831	16	0	5.831	15	0	6.606	15	0	7.034	1	3	5.694	14	0	3.583	14	0	3.583		
2	1	5.723	7.594	16	1	7.594	15	1	7.796	15	1	4.319	1	4	6.810	14	1	4.993	14	1	4.993		
2	2	7.225	9.117	16	2	9.117	15	2	2.626	15	2	5.874	2	0	2.037	14	2	6.206	14	2	6.206		
2	3	8.619	5.979	17	0	5.979	16	0	4.113	16	0	7.187	2	1	3.387	15	0	3.735	15	0	3.735		
3	0	4.087	7.747	17	1	7.747	16	1	2.543	16	1	6.032	2	2	4.605	15	1	5.150	15	1	5.150		
3	1	5.816	9.272	17	2	9.272	16	2	6.671	16	2	4.468	2	3	5.759	15	2	6.368	15	2	6.368		
3	2	7.319	6.128	18	0	6.128	17	0	2.719	17	0	4.617	3	0	2.129	16	0	3.884	16	0	3.884		
3	3	8.712	7.901	18	1	7.901	17	1	5.525	17	1	6.190	3	1	3.480	16	1	5.307	16	1	5.307		
4	0	4.194	9.427	18	2	9.427	18	0	4.206	18	0	7.496	3	2	4.698	17	0	4.035	17	0	4.035		
4	1	5.924	6.278	19	0	6.278	18	1	6.765	18	1	4.768	3	3	5.854	17	1	5.464	17	1	5.464		
4	2	7.427	8.056	19	1	8.056	18	2	2.826	18	2	7.651	4	0	2.237	18	0	4.186	18	0	4.186		
4	3	8.821	9.583	19	2	9.583	19	0	5.635	19	0	4.919	4	1	3.589	18	1	5.623	18	1	5.623		
5	0	4.311	6.429	20	0	6.429	19	1	2.719	19	1	6.506	4	2	4.808	19	0	4.339	19	0	4.339		
5	1	6.042	8.212	20	1	8.212	20	0	2.943	20	0	5.072	4	3	5.964	19	1	5.780	19	1	5.780		
5	2	7.546	9.381	21	0	9.381	21	0	4.433	21	0	6.824	5	0	2.354	20	0	4.491	20	0	4.491		
5	3	8.941	6.741	21	1	6.741	21	1	5.754	21	1	5.532	5	1	3.708	20	1	5.938	20	1	5.938		
6	0	4.435	7.411	22	0	7.411	22	0	6.995	22	0	7.123	6	0	2.478	21	0	4.645	21	0	4.645		
6	1	6.167	8.538	22	1	8.538	22	1	3.067	22	1	6.982	6	1	3.834	21	1	6.096	21	1	6.096		
6	2	7.672	9.896	23	0	9.896	23	0	4.559	23	0	7.141	6	2	5.056	22	0	4.799	22	0	4.799		
6	3	9.067	6.896	23	1	6.896	24	0	6.013	24	0	5.687	6	3	6.216	22	1	6.254	22	1	6.254		
7	0	4.564	7.050	24	0	7.050	24	1	7.123	24	1	7.300	7	0	2.606	23	0	4.954	23	0	4.954		
7	1	6.297	8.855	24	1	8.855	25	0	3.196	25	0	5.843	7	1	3.966	23	1	6.411	23	1	6.411		
7	2	7.803	7.206	25	0	7.206	25	1	4.690	25	1	7.141	7	2	5.189	24	0	5.109	24	0	5.109		
7	3	9.199	9.014	25	1	9.014	26	0	6.013	26	0	5.687	7	3	6.347	24	1	6.569	24	1	6.569		
8	0	4.696	7.362	26	0	7.362	26	1	7.256	26	1	7.300	8	0	2.739	25	0	5.264	25	0	5.264		
8	1	6.435	9.173	26	1	9.173	27	0	3.328	27	0	5.843	8	1	3.966	25	1	6.727	25	1	6.727		
8	2	7.939	7.519	27	0	7.519	27	1	4.825	27	1	7.459	8	2	5.327	26	0	5.419	26	0	5.419		
8	3	9.336	9.333	27	1	9.333	28	0	6.150	28	0	7.617	8	3	6.485	26	1	6.885	26	1	6.885		
9	0	4.831	7.675	28	0	7.675	28	1	7.394	28	1	6.154	9	0	2.874	27	0	5.575	27	0	5.575		
9	1	6.569	9.492	28	1	9.492	29	0	3.463	29	0	7.776	9	1	4.241	28	0	5.731	28	0	5.731		
9	2	8.078	7.833	29	0	7.833	29	1	4.964	29	1	6.310	9	2	5.469	29	0	5.887	29	0	5.887		
9	3	9.476	7.991	30	0	7.991	30	1	6.290	30	1	6.467	9	3	6.627	30	0	6.044	30	0	6.044		
10	0	4.969	8.148	31	0	8.148	31	1	7.536	31	1	6.624	10	0	3.012	31	0	6.201	31	0	6.201		
10	1	6.709	8.307	32	0	8.307	32	1	3.601	32	1	7.096	10	1	4.384	32	0	6.357	32	0	6.357		
10	2	8.223	8.465	33	0	8.465	33	1	5.105	33	1	7.681	10	2	5.610	33	0	6.514	33	0	6.514		
11	0	5.109	8.624	34	0	8.624	34	1	5.105	34	1	7.096	11	0	3.741	34	0	6.671	34	0	6.671		
11	1	6.852	8.783	35	0	8.783	35	1	7.681	35	1	7.253	11	1	6.772	35	0	6.828	35	0	6.828		
11	2	8.364	8.942	36	0	8.942	36	1	3.741	36	1	7.411	11	2	4.529	36	0	6.986	36	0	6.986		
12	0	5.251	9.101	37	0	9.101	37	1	5.249	37	1	7.569	12	0	3.152	37	0						
12	1	6.997	9.260	38	0	9.260	38	1	6.579	38	1	7.727	12	1	1								
12	2	8.510	9.420	39	0	9.420	39	1	7.829	39	1												
13	0	5.594	9.580	40	0	9.580	40	1															

Table 1. (Continued)

a1 = 0.2    alpha = 0.01				a1 = 0.2    alpha = 0.05				a1 = 0.2    alpha = 0.10			
x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)
0	0	3.684	10	2	8.377	0	0	2.397	9	3	7.664
0	1	5.311	10	3	9.716	0	1	3.795	10	0	3.982
0	2	6.725	11	0	5.469	0	2	5.037	10	1	5.440
0	3	8.036	11	1	7.153	0	3	6.203	10	2	6.697
0	4	9.284	11	2	8.590	0	4	7.323	10	3	7.873
1	0	3.714	11	3	9.938	0	5	8.410	11	0	4.187
1	1	5.341	12	0	5.676	1	0	2.427	11	1	5.652
1	2	6.755	12	1	7.381	1	1	3.825	11	2	6.908
1	3	8.067	12	2	8.805	1	2	5.067	11	3	8.084
1	4	9.314	12	3	10.155	1	3	6.233	12	0	4.395
2	0	3.821	13	0	5.885	1	4	7.353	12	1	5.665
2	1	5.449	13	1	7.598	1	5	8.441	12	2	7.120
2	2	6.864	13	2	9.020	2	0	2.534	12	3	8.295
2	3	8.176	14	0	6.096	2	1	3.934	13	0	4.605
2	4	9.424	14	1	7.815	2	2	5.176	13	1	6.158
3	0	3.965	14	2	9.236	2	3	6.343	13	2	7.332
3	1	5.595	15	0	6.308	2	4	7.464	13	3	8.506
3	2	7.011	15	1	8.034	2	5	8.552	14	0	4.816
3	3	8.324	15	2	9.452	3	0	2.677	14	1	6.320
3	4	9.573	16	0	6.522	3	1	4.081	14	2	7.544
4	0	4.128	16	1	8.252	3	2	5.325	15	0	5.072
4	1	5.761	16	2	9.669	3	3	6.493	15	1	6.536
4	2	7.179	17	0	6.776	3	4	7.614	15	2	7.757
4	3	8.494	17	1	8.471	4	0	4.248	16	0	5.288
4	4	9.744	17	2	9.845	4	1	5.496	16	1	6.752
5	0	4.302	18	0	6.996	4	2	6.664	16	2	7.970
5	1	5.940	18	1	8.691	4	3	7.787	17	0	5.506
5	2	7.361	18	2	10.060	4	4	8.915	17	1	6.967
5	3	8.678	19	0	7.216	5	0	3.015	17	2	8.183
5	4	9.926	19	1	8.910	5	1	4.430	18	0	5.723
6	0	4.485	19	2	10.276	5	2	5.681	18	1	7.183
6	1	6.129	20	0	7.436	5	3	6.850	18	2	8.397
6	2	7.552	20	1	9.129	5	4	7.975	19	0	5.941
6	3	8.872	21	0	7.657	6	0	3.198	19	1	7.399
6	4	10.118	21	1	9.349	6	1	4.640	20	0	6.104
7	0	4.674	22	0	7.877	6	2	5.890	20	1	7.614
7	1	6.339	22	1	9.569	6	3	7.044	21	0	6.376
7	2	7.750	23	0	8.098	6	4	8.172	21	1	7.830
7	3	9.075	23	1	9.886	7	0	3.366	22	0	6.594
7	4	10.316	24	0	8.319	7	1	4.855	22	1	8.046
8	0	4.867	24	1	10.103	7	2	6.066	23	0	6.812
8	1	6.557	25	0	8.540	7	3	7.244	23	1	8.262
8	2	7.954	26	0	8.761	7	4	8.375	24	0	7.029
8	3	9.285	27	0	9.982	8	0	3.579	24	1	8.477
9	0	5.064	28	0	9.203	8	1	5.019	25	0	7.247
9	1	6.729	29	0	9.425	8	2	6.269	26	0	7.465
9	2	8.161	30	0	9.646	8	3	7.449	27	0	7.683
9	3	9.500	31	0	9.802	9	0	3.776	28	0	7.901
10	0	5.264	32	0	10.023	9	1	5.230	29	0	8.119
10	1	6.940	33	0	10.244	9	2	6.474	30	0	8.337

Table 1. (Continued)

a1 = 0.25				alpha = 0.01				a1 = 0.25				alpha = 0.05				a1 = 0.25				alpha = 0.10			
x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)
0	0	3.454	8	4	10.491	0	0	2.247	8	1	5.307	0	0	1.727	8	0	3.441						
0	1	4.979	9	0	5.406	0	1	3.558	8	2	6.432	0	1	2.917	8	1	4.715						
0	2	6.304	9	1	7.041	0	2	4.722	8	3	7.527	0	2	3.992	8	2	5.756						
0	3	7.534	9	2	8.345	0	3	5.815	8	4	8.680	0	3	5.011	8	3	6.827						
0	4	8.704	9	3	9.581	0	4	6.865	9	0	4.210	0	4	5.995	8	4	7.741						
0	5	9.831	9	4	10.815	0	5	7.885	9	1	5.686	0	5	6.956	9	0	3.716						
1	0	3.508	10	0	5.718	0	6	8.882	9	2	6.809	0	6	7.899	9	1	4.986						
1	1	5.033	10	1	7.325	1	0	2.301	9	3	7.907	1	0	1.781	9	2	6.022						
1	2	6.359	10	2	8.624	1	1	3.613	9	4	8.950	1	1	2.972	9	3	7.028						
1	3	7.589	10	3	9.857	1	2	4.777	10	0	4.486	1	2	4.047	9	4	8.005						
1	4	8.762	11	0	6.003	1	3	5.871	10	1	5.856	1	3	5.066	10	0	3.991						
1	5	9.887	11	1	7.609	1	4	6.921	10	2	7.087	1	4	6.051	10	1	5.261						
2	0	3.671	11	2	8.902	1	5	7.941	10	3	8.181	1	5	7.012	10	2	6.292						
2	1	5.200	11	3	10.133	1	6	8.938	11	0	4.830	1	6	7.955	10	3	7.294						
2	2	6.528	12	0	6.288	2	0	2.464	11	1	6.243	2	0	1.944	11	0	4.268						
2	3	7.759	12	1	7.893	2	1	3.781	11	2	7.360	2	1	3.142	11	1	5.533						
2	4	8.931	12	2	9.183	2	2	4.948	11	3	8.455	2	2	4.221	11	2	6.490						
2	5	10.060	12	3	10.410	2	3	6.044	12	0	5.111	2	3	5.242	11	3	7.559						
3	0	3.878	13	0	6.574	2	4	7.096	12	1	6.524	2	4	6.233	12	0	4.545						
3	1	5.435	13	1	8.178	2	5	8.118	12	2	7.633	2	5	7.186	12	1	5.805						
3	2	6.758	13	2	9.463	3	0	2.671	12	3	8.594	2	6	8.130	12	2	6.756						
3	3	7.981	13	3	10.687	3	1	4.001	13	0	5.392	3	0	2.151	12	3	7.825						
3	4	9.155	14	0	6.861	3	2	5.172	13	1	6.884	3	1	3.362	13	0	4.823						
3	5	10.288	14	1	8.462	3	3	6.274	13	2	7.781	3	2	4.450	13	1	6.077						
4	0	4.108	14	2	9.748	3	4	7.333	13	3	8.863	3	3	5.489	13	2	7.095						
4	1	5.656	15	0	7.148	3	5	8.368	14	0	5.607	3	4	6.444	13	3	8.091						
4	2	6.988	15	1	8.747	4	0	2.901	14	1	6.961	3	5	7.407	14	0	5.101						
4	3	8.228	15	2	10.019	4	1	4.267	14	2	8.058	4	0	2.381	14	1	6.350						
4	4	9.407	16	0	7.435	4	2	5.422	15	0	5.954	4	1	3.607	14	2	7.363						
4	5	10.545	16	1	9.031	4	3	6.537	15	1	7.237	4	2	4.668	14	3	8.230						
5	0	4.352	16	2	10.299	4	4	7.599	15	2	8.331	4	3	5.692	15	0	5.379						
5	1	5.914	17	0	7.723	4	5	8.569	16	0	6.174	4	4	6.681	15	1	6.622						
5	2	7.244	17	1	9.319	5	0	3.145	16	1	7.514	5	0	2.625	15	2	7.712						
5	3	8.489	17	2	10.579	5	1	4.549	16	2	8.729	5	1	3.865	16	0	5.658						
5	4	9.672	18	0	8.010	5	2	5.639	17	0	6.458	5	2	4.913	16	1	6.895						
5	5	10.754	18	1	9.603	5	3	6.738	17	1	7.869	5	3	5.938	16	2	7.952						
6	0	4.605	18	2	10.860	5	4	7.793	17	2	9.003	5	4	6.928	17	0	5.937						
6	1	6.182	19	0	8.298	5	5	8.815	18	0	8.143	5	5	7.894	17	1	7.167						
6	2	7.507	19	1	9.888	6	0	3.398	18	1	8.143	6	0	2.878	18	0	6.216						
6	3	8.757	20	0	8.586	6	1	4.755	19	0	7.079	6	1	4.134	18	1	7.440						
6	4	9.944	20	1	10.173	6	2	5.961	19	1	8.418	6	2	5.173	19	0	6.498						
7	0	4.864	21	0	8.874	6	3	7.000	20	0	7.369	6	3	6.197	19	1	7.590						
7	1	6.459	21	1	10.458	6	4	8.052	20	1	8.628	6	4	7.215	20	0	6.777						
7	2	7.792	22	0	9.162	6	5	9.075	21	0	9.066	6	5	8.175	21	0	7.055						
7	3	9.036	22	1	10.743	7	0	3.665	21	1	8.906	7	0	3.168	21	1	8.133						
7	4	10.217	23	0	9.450	7	1	5.035	22	0	7.932	7	1	4.403	22	0	7.336						
8	0	5.133	24	0	9.738	7	2	6.166	23	0	8.213	7	2	5.434	23	0	7.682						
8	1	6.738	25	0	10.036	7	3	7.262	24	0	8.495	7	3	6.559	24	0	7.985						
8	2	8.068	26	0	10.325	7	4	8.312	25	0	8.777	7	4	7.478	25	0	8.232						
8	3	9.307	27	0	10.613	8	0	3.936	26	0	9.058	8	0										

Table 1. (Continued)

a1 = 0.3					alpha = 0.01					a1 = 0.3					alpha = 0.05					a1 = 0.3					alpha = 0.10				
x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)			
0	0	3.224	7	2	7.907	0	0	2.097	7	1	5.319	0	0	1.612	6	5	8.102	0	0	1.612	6	5	8.102	0	0	1.612			
0	1	4.647	7	3	9.143	0	1	3.321	7	2	6.465	0	1	2.723	7	0	3.581	0	1	2.723	7	0	3.581	0	1	2.723			
0	2	5.884	7	4	10.206	0	2	4.407	7	3	7.427	0	2	3.726	7	1	4.761	0	2	3.726	7	1	4.761	0	2	3.726			
0	3	7.032	7	5	11.097	0	3	5.428	7	4	8.440	0	3	4.676	7	2	5.773	0	3	4.676	7	2	5.773	0	3	4.676			
0	4	8.123	8	0	5.549	0	4	6.407	7	5	9.386	0	4	5.597	7	3	6.639	0	4	5.597	7	3	6.639	0	4	5.597			
0	5	9.176	8	1	7.104	0	5	7.359	8	0	4.446	0	5	6.492	7	4	7.540	0	5	6.492	7	4	7.540	0	5	6.492			
0	6	10.199	8	2	8.257	0	6	8.290	8	1	5.751	0	6	7.373	7	5	8.260	0	6	7.373	7	5	8.260	0	6	7.373			
0	7	11.200	8	3	9.476	0	7	9.204	8	2	6.810	0	7	8.240	8	0	3.929	0	7	8.240	8	0	3.929	0	7	8.240			
1	0	3.315	8	4	10.545	1	0	2.189	8	3	7.667	1	0	1.703	8	1	5.101	1	0	1.703	8	1	5.101	1	0	1.703			
1	1	4.741	9	0	5.910	1	1	3.416	8	4	8.620	1	1	2.818	8	2	6.109	1	1	2.818	8	2	6.109	1	1	2.818			
1	2	5.979	9	1	7.465	1	2	4.504	9	0	4.801	1	2	3.823	8	3	6.965	1	2	3.823	8	3	6.965	1	2	3.823			
1	3	7.128	9	2	8.607	1	3	5.525	9	1	6.006	1	3	4.776	8	4	7.862	1	3	4.776	8	4	7.862	1	3	4.776			
1	4	8.220	9	3	9.816	1	4	6.506	9	2	6.987	1	4	5.696	9	0	4.279	1	4	5.696	9	0	4.279	1	4	5.696			
1	5	9.274	9	4	10.885	1	5	7.459	9	3	8.164	1	5	6.594	9	1	5.284	1	5	6.594	9	1	5.284	1	5	6.594			
1	6	10.298	10	0	6.273	1	6	8.390	9	4	9.253	1	6	7.475	9	2	6.413	1	6	7.475	9	2	6.413	1	6	7.475			
2	0	3.552	10	1	7.727	2	0	3.305	10	0	5.155	2	0	1.940	9	3	7.296	2	0	1.940	9	3	7.296	2	0	1.940			
2	1	4.992	10	2	8.959	2	1	2.425	10	1	6.354	2	1	3.074	10	0	4.629	2	1	3.074	10	0	4.629	2	1	3.074			
2	2	6.232	10	3	9.989	2	2	3.666	10	2	7.487	2	2	4.096	10	1	5.625	2	2	4.096	10	1	5.625	2	2	4.096			
2	3	7.386	10	4	11.248	2	3	4.765	10	3	8.341	2	3	5.060	10	2	6.745	2	3	5.060	10	2	6.745	2	3	5.060			
2	4	8.484	11	0	6.637	2	4	5.798	10	4	9.284	2	4	5.932	10	3	7.612	2	4	5.932	10	3	7.612	2	4	5.932			
2	5	9.543	11	1	8.081	2	5	6.785	11	0	5.510	2	5	6.830	10	4	8.508	2	5	6.830	10	4	8.508	2	5	6.830			
2	6	10.573	11	2	9.311	2	6	7.695	11	1	6.699	2	6	7.711	11	0	4.982	2	6	7.711	11	0	4.982	2	6	7.711			
3	0	3.840	11	3	10.332	2	6	8.627	11	2	7.825	2	6	8.579	11	1	5.950	2	7	8.579	11	1	5.950	2	7	8.579			
3	1	5.310	12	0	7.001	3	0	2.714	11	3	8.772	2	7	9.228	11	2	7.076	3	0	2.714	11	2	7.076	3	0	2.714			
3	2	6.538	12	1	8.435	3	1	3.970	12	0	5.865	3	1	3.384	11	3	7.937	3	1	3.384	11	3	7.937	3	1	3.384			
3	3	7.699	12	2	9.665	3	2	5.082	12	1	7.108	3	2	4.340	11	4	8.672	3	2	4.340	11	4	8.672	3	2	4.340			
3	4	8.804	12	3	10.675	3	3	6.044	12	2	7.999	3	3	5.294	12	0	5.399	3	3	5.294	12	0	5.399	3	3	5.294			
3	5	9.870	13	0	7.367	3	4	7.026	12	3	9.104	3	4	6.212	12	1	6.283	3	4	6.212	12	1	6.283	3	4	6.212			
3	6	10.909	13	1	8.788	3	5	8.065	13	0	6.121	3	5	7.111	12	2	7.407	3	5	7.111	12	2	7.407	3	5	7.111			
4	0	4.154	13	2	10.089	3	6	8.907	13	1	7.289	3	6	8.502	13	0	5.739	3	6	8.502	13	0	5.739	3	6	8.502			
4	1	5.663	13	3	11.018	4	0	3.027	13	2	8.436	4	0	6.577	13	1	6.617	4	0	3.027	13	1	6.617	4	0	3.027			
4	2	6.865	14	0	7.826	4	1	4.294	13	3	9.436	4	1	5.412	14	0	6.179	4	1	4.294	14	0	6.179	4	1	4.294			
4	3	8.028	14	1	9.133	4	2	5.412	14	0	6.577	4	2	5.590	14	0	6.079	4	2	5.412	14	0	6.079	4	2	5.412			
4	4	9.042	14	2	10.438	4	3	6.341	14	1	7.629	4	3	5.951	14	1	6.951	4	3	6.341	14	1	6.951	4	3	6.341			
4	5	10.097	15	0	8.186	4	4	7.322	14	2	8.841	4	4	6.517	14	2	8.068	4	4	7.322	14	2	8.068	4	4	7.322			
4	6	11.212	15	1	9.496	4	5	8.275	15	0	6.934	4	5	7.415	15	0	6.446	4	5	8.275	15	0	6.446	4	5	8.275			
5	0	4.483	15	2	10.785	4	6	9.213	15	1	8.131	4	6	8.425	15	1	7.284	4	6	9.213	15	1	7.284	4	6	9.213			
5	1	6.024	16	0	8.547	5	0	3.360	15	2	7.179	4	6	8.425	15	1	7.284	5	0	3.360	15	1	7.284	5	0	3.360			
5	2	7.214	16	1	9.850	5	1	4.637	16	0	7.387	5	1	4.047	15	2	8.399	5	1	4.637	16	0	8.399	5	1	4.637			
5	3	8.371	16	2	11.131	5	2	5.658	16	1	8.310	5	2	4.962	16	1	7.530	5	2	5.658	16	1	7.530	5	2	5.658			
5	4	9.382	17	0	8.909	5	3	6.670	16	2	9.518	5	3	5.992	16	2	8.573	5	3	6.670	16	2	8.573	5	3	6.670			
5	5	10.433	17	1	10.203	5	4	7.777	17	0	7.735	5	4	6.898	16	3	7.963	5	4	7.777	17	0	7.963	5	4	7.777			
6	0	4.837	18	0	9.270	6	0	3.742	17	1	8.652	5	5	7.787	17	0	7.118	6	0	3.742	17	1	7.118	6	0	3.742			
6	1	6.384	18	1	10.557	6	1	4.977	18	0	8.008	6	1	4.225	17	1	7.963	6	1	4.977	18	0	7.963	6	1	4.977			
6	2	7.559	19	0	9.631	6	2	6.129	19	0	8.426	6	2	3.235	18	1	8.296	6	2	6.129	19	0	8.296	6	2	6.129			
6	3	8.712	19	1	10.911	6	3	7.148	19	1	9.333	6	3	4.427	18	2	8.455	6	3	7.148	19	1	8.455	6	3	7.148			
6	4	9.714	20	0	9.994	6	4	7.969	20	0	8.770	6	4	5.437	19	2	9.791	6	4	7.969	20	0	9.791	6	4	7.969			
6	5	10.763	21	0	10.356	6	5	9.056	20	1	9.117	6	5	6.311	19	3	8.128	6	5	9.056	20	1	8.128	6	5	9.056			
7	0	5.192	22	0	10.718	6	6	4.094	21	0	9.462	6	6	7.215	20	0	8.465	7	0	5.192	22	0	8.465	7	0	5.192			
7	1	6.744	23	0	11.080	7	0	4.094	22	0	9.462	7	1	6.744	21	1	8.465	7	1	6.744	23	0	8.465	7	1	6.744			

Table 1. (Continued)

a1 = 0.35    alpha = 0.01				a1 = 0.35    alpha = 0.05				a1 = 0.35    alpha = 0.10			
x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)	x1	x2	t(x1,x2)
0	0	2.993	6	2	7.650	0	0	1.947	6	0	3.687
0	1	4.315	6	3	8.678	0	1	3.084	6	1	4.547
0	2	5.464	6	4	9.681	0	2	4.092	6	2	5.636
0	3	6.529	6	5	10.649	0	3	5.040	6	3	6.490
0	4	7.543	6	6	11.583	0	4	5.950	6	4	7.323
0	5	8.520	7	0	5.763	0	5	6.834	6	5	7.950
0	6	9.471	7	1	6.965	0	6	7.698	6	6	8.949
0	7	10.400	7	2	8.082	0	7	8.546	7	0	4.120
0	8	11.312	7	3	9.102	0	8	9.383	7	1	4.947
1	0	3.146	7	4	10.094	1	0	2.100	7	2	5.962
1	1	4.478	7	5	11.059	1	1	3.243	7	3	6.946
1	2	5.672	8	0	6.211	1	2	4.256	7	4	7.761
1	3	6.692	8	1	7.396	1	3	5.207	7	5	8.333
1	4	7.708	8	2	8.510	1	4	6.119	8	0	4.641
1	5	8.687	8	3	9.520	1	5	7.005	8	1	5.348
1	6	9.639	8	4	10.508	1	6	7.871	8	2	6.343
1	7	10.569	8	5	11.299	1	7	8.721	8	3	7.071
1	8	11.482	9	0	6.659	9	0	5.559	8	4	8.096
2	0	3.485	9	1	7.828	2	0	2.439	8	5	8.718
2	1	4.871	9	2	8.937	2	1	3.587	9	0	5.048
2	2	5.971	9	3	9.938	2	2	4.603	9	1	5.749
2	3	7.043	9	4	10.923	2	3	5.557	9	2	6.749
2	4	8.056	10	0	7.108	2	4	6.471	9	3	7.462
2	5	9.037	10	1	8.260	2	5	7.359	9	4	8.483
2	6	9.990	10	2	9.345	2	6	8.123	10	0	5.454
2	7	10.921	10	3	10.356	2	7	8.971	10	1	6.149
3	0	3.881	10	4	11.470	3	0	2.834	10	2	7.210
3	1	5.203	11	0	7.557	3	1	3.971	10	3	7.852
3	2	6.361	11	1	8.721	3	2	4.984	10	4	8.870
3	3	7.440	11	2	9.768	3	3	5.818	11	0	5.860
3	4	8.439	11	3	10.774	3	4	6.724	11	1	6.548
3	5	9.419	12	0	8.007	3	5	7.743	11	2	7.603
3	6	10.373	12	1	9.151	3	6	8.608	11	3	8.243
3	7	11.179	12	2	10.191	3	7	9.312	12	0	6.266
4	0	4.304	12	3	11.191	4	0	3.301	12	1	6.924
4	1	5.615	13	0	8.457	4	1	4.402	12	2	7.996
4	2	6.803	13	1	9.580	4	2	5.399	12	3	8.633
4	3	7.869	13	2	10.613	4	3	6.342	13	0	6.671
4	4	8.859	14	0	8.909	4	4	7.140	13	1	7.344
4	5	9.834	14	1	10.008	4	5	8.241	13	2	8.389
4	6	10.783	14	2	11.034	4	6	9.078	14	0	7.078
5	0	4.758	15	0	9.365	4	7	9.718	14	1	7.736
5	1	6.103	15	1	10.437	5	0	3.736	14	2	8.781
5	2	7.226	15	2	11.437	5	1	4.817	15	0	7.480
5	3	8.291	16	0	9.811	5	2	5.998	15	1	8.137
5	4	9.267	16	1	10.864	5	3	6.860	16	0	7.881
5	5	10.239	17	0	10.258	5	4	7.528	16	1	8.532
5	6	11.374	17	1	11.291	5	5	8.401	17	0	8.282
6	0	5.313	18	0	10.703	5	6	9.460	17	1	8.927
6	1	6.533	19	0	11.146	6	0	4.174	18	0	8.682

Table 1. (Continued)

a1 = 0.4				alpha = 0.01				a1 = 0.4				alpha = 0.05				a1 = 0.4				alpha = 0.10			
x1	x2	t(x1,x2)		x1	x2	t(x1,x2)		x1	x2	t(x1,x2)		x1	x2	t(x1,x2)		x1	x2	t(x1,x2)		x1	x2	t(x1,x2)	
0	0	2.763		5	4	9.059		0	0	1.797		5	3	6.770		0	0	1.382		5	3	6.053	
0	1	3.983		5	5	9.926		0	1	2.846		5	4	7.580		0	1	2.334		5	4	6.792	
0	2	5.044		5	6	10.889		0	2	3.778		5	5	8.373		0	2	3.193		5	5	7.349	
0	3	6.027		6	0	5.902		0	3	4.652		5	6	9.153		0	3	4.008		5	6	8.414	
0	4	6.963		6	1	6.748		0	4	5.492		5	7	9.913		0	4	4.796		5	7	8.821	
0	5	7.865		6	2	7.943		0	5	6.308		6	0	4.737		0	5	5.565		6	0	4.213	
0	6	8.742		6	3	8.651		0	6	7.114		6	1	5.576		0	6	6.319		6	1	4.951	
0	7	9.600		6	4	9.763		0	7	7.889		6	2	6.421		0	7	7.063		6	2	5.739	
0	8	10.442		6	5	10.521		0	8	8.661		6	3	7.241		0	8	7.797		6	3	6.305	
0	9	11.270		6	6	11.370		0	9	9.423		6	4	8.047		0	9	8.524		6	4	7.564	
1	0	3.022		7	0	6.442		1	0	2.057		6	5	8.836		1	0	1.641		6	5	8.016	
1	1	4.270		7	1	7.258		1	1	3.080		6	6	9.577		1	1	2.563		6	6	9.033	
1	2	5.368		7	2	8.456		1	2	4.103		7	0	5.259		1	2	3.416		7	0	4.708	
1	3	6.258		7	3	9.370		1	3	4.876		7	1	6.065		1	3	4.228		7	1	5.427	
1	4	7.190		7	4	10.260		1	4	5.713		7	2	6.899		1	4	5.014		7	2	6.446	
1	5	8.090		7	5	11.006		1	5	6.528		7	3	7.712		1	5	5.782		7	3	6.973	
1	6	8.966		7	6	11.703		1	6	7.324		7	4	8.523		1	6	6.535		7	4	7.796	
1	7	9.822		8	0	6.980		1	7	8.107		7	5	9.301		1	7	7.277		7	5	8.251	
1	8	10.663		8	1	7.797		1	8	8.878		8	0	5.774		1	8	8.011		8	0	5.198	
1	9	11.491		8	2	8.967		1	9	9.640		8	1	6.552		1	9	8.737		8	1	5.891	
2	0	3.514		8	3	9.646		2	0	2.549		8	2	7.377		2	0	2.133		8	2	6.833	
2	1	4.784		8	4	10.635		2	1	3.503		8	3	8.183		2	1	2.974		8	3	7.216	
2	2	5.843		8	5	11.490		2	2	4.415		8	4	8.983		2	2	3.818		8	4	8.467	
2	3	6.675		9	0	7.519		2	3	5.282		8	5	9.751		2	3	4.626		8	5	8.702	
2	4	7.604		9	1	8.317		2	4	6.118		9	0	6.282		2	4	5.409		9	0	5.682	
2	5	8.502		9	2	9.248		2	5	6.932		9	1	7.037		2	5	6.175		9	1	6.148	
2	6	9.376		9	3	10.145		2	6	7.728		9	2	7.854		2	6	6.927		9	2	7.386	
2	7	10.227		9	4	11.126		2	7	8.502		9	3	8.653		2	7	7.668		9	3	7.674	
2	8	11.072		10	0	8.058		2	8	9.272		9	4	9.615		2	8	8.358		9	4	8.878	
3	0	4.077		10	1	8.835		2	9	10.033		10	0	6.785		2	9	9.127		9	5	9.153	
3	1	5.169		10	2	9.986		3	0	3.158		10	1	7.521		3	0	2.681		10	0	6.282	
3	2	6.394		10	3	10.751		3	1	3.982		10	2	8.330		3	1	3.532		10	1	6.615	
3	3	7.287		10	4	11.784		3	2	5.086		10	3	9.123		3	2	4.352		10	2	7.889	
3	4	8.193		11	0	8.596		3	3	5.831		10	4	9.985		3	3	5.144		10	3	8.131	
3	5	9.169		11	1	9.476		3	4	6.651		11	0	7.283		3	4	5.917		11	0	6.752	
3	6	10.038		11	2	10.370		3	5	7.452		11	1	8.002		3	5	6.672		11	1	7.080	
3	7	10.780		11	3	11.248		3	6	8.238		11	2	8.804		3	6	7.485		11	2	8.301	
3	8	11.611		12	0	9.128		3	7	9.013		11	3	9.397		3	7	8.154		11	3	8.587	
4	0	4.655		12	1	9.863		3	8	9.778		12	0	7.777		3	8	8.948		12	0	7.236	
4	1	5.687		12	2	10.868		4	0	3.679		12	1	8.482		4	1	3.197		12	1	7.539	
4	2	6.909		12	3	11.634		4	1	4.581		12	2	9.524		4	2	3.996		12	2	8.758	
4	3	7.760		13	0	9.656		4	2	5.450		12	3	9.868		4	3	4.814		12	3	9.042	
4	4	8.773		13	1	10.489		4	3	6.289		13	0	8.267		4	4	5.598		13	0	7.701	
4	5	9.539		13	2	11.365		4	4	7.104		13	1	8.960		4	5	6.510		13	1	8.005	
4	6	10.397		14	0	10.178		4	5	7.911		13	2	10.059		4	6	7.117		13	2	9.213	
4	7	11.239		14	1	10.992		4	6	8.694		14	0	8.755		4	7	7.865		14	0	8.163	
5	0	5.325		15	0	10.697		4	7	9.461		14	1	9.229		4	8	8.593		15	0	8.460	
5	1	6.210		15	1	11.497		5	0	4.209		15	0	9.357		5	1	3.710		15	0	8.624	
5	2	7.628		16	0	11.212		5	1	4.971		15	1	9.703		5	2	4.478		15	1	8.925	
5	3	8.277		17	0	11.724		5	2	5.942		16	0	9.833		5	3	5.276		16	0	9.084	



Table 1. (Continued)

a1 = 0.45				alpha = 0.01				a1 = 0.45				alpha = 0.05				a1 = 0.45				alpha = 0.10						
x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)	x1	x2	t(x1,x2)	t(x1,x2)			
0	0	2.533	7	10.825	4	7	10.825	0	0	1.648	6	8.441	4	6	8.441	0	0	1.266	4	6	7.795	4	6	7.795		
0	1	3.651	4	11.574	4	7	11.574	0	1	2.609	4	7	9.139	4	7	9.139	0	1	2.139	4	7	8.490	4	7	8.490	
0	2	4.623	5	0	5.899	4	8	9.829	0	2	3.490	4	8	9.829	4	8	9.829	0	2	2.964	4	8	9.116	4	8	9.116
0	3	5.525	5	1	6.666	4	8	10.809	0	3	4.264	5	0	4.809	5	0	4.809	0	3	3.674	5	0	4.233	5	0	4.233
0	4	6.382	5	2	7.491	4	9	11.574	0	4	5.034	5	1	5.270	5	1	5.270	0	4	4.419	5	1	4.540	5	1	4.540
0	5	7.217	5	3	8.300	4	10	12.349	0	5	5.782	5	2	6.124	5	2	6.124	0	5	5.101	5	2	5.470	5	2	5.470
0	6	8.014	5	4	9.107	4	11	13.129	0	6	6.513	5	3	6.848	5	3	6.848	0	6	5.793	5	3	6.152	5	3	6.152
0	7	8.800	5	5	9.870	4	12	13.914	0	7	7.232	5	4	7.591	5	4	7.591	0	7	6.474	5	4	6.856	5	4	6.856
0	8	9.571	5	6	10.634	4	13	14.709	0	8	7.939	5	5	8.283	5	5	8.283	0	8	7.147	5	5	7.505	5	5	7.505
0	9	10.331	5	7	11.387	4	14	15.504	0	9	8.656	5	6	8.984	5	6	8.984	0	9	7.813	5	6	8.173	5	6	8.173
0	10	11.092	6	0	6.557	5	0	6.557	0	10	9.329	5	7	9.677	5	7	9.677	0	10	8.474	5	7	8.834	5	7	8.834
0	11	11.833	6	1	7.129	5	1	7.129	0	11	10.033	6	0	5.397	6	0	5.397	0	11	9.129	6	0	4.791	6	0	4.791
1	0	3.021	6	2	8.099	5	2	8.099	1	0	2.135	6	1	5.843	6	1	5.843	1	0	1.750	6	1	5.198	6	1	5.198
1	1	4.073	6	3	8.894	5	3	8.894	1	1	3.011	6	2	6.692	6	2	6.692	1	1	2.528	6	2	6.010	6	2	6.010
1	2	5.020	6	4	9.693	5	4	9.693	1	2	3.839	6	3	7.409	6	3	7.409	1	2	3.290	6	3	6.685	6	3	6.685
1	3	5.908	6	5	10.444	5	5	10.444	1	3	4.626	6	4	8.141	6	4	8.141	1	3	4.024	6	4	7.385	6	4	7.385
1	4	6.757	6	6	11.180	5	6	11.180	1	4	5.391	6	5	8.672	6	5	8.672	1	4	4.742	6	5	8.027	6	5	8.027
1	5	7.578	6	7	11.929	5	7	11.929	1	5	6.134	6	6	9.521	6	6	9.521	1	5	5.437	6	6	8.691	6	6	8.691
1	6	8.377	6	8	12.678	5	8	12.678	1	6	6.861	6	7	10.055	6	7	10.055	1	6	6.124	6	7	5.344	6	7	5.344
1	7	9.160	7	1	7.803	6	1	7.803	1	7	7.569	7	0	5.977	7	0	5.977	1	7	6.802	7	0	5.703	7	0	5.703
1	8	9.928	7	2	8.721	6	2	8.721	1	8	8.273	7	1	6.371	7	1	6.371	1	8	7.472	7	1	6.547	7	1	6.547
1	9	10.685	7	3	9.506	6	3	9.506	1	9	8.969	7	2	7.252	7	2	7.252	1	9	8.136	7	2	7.055	7	2	7.055
1	10	11.432	7	4	10.286	6	4	10.286	1	10	9.659	7	3	7.961	7	3	7.961	1	10	8.794	7	3	7.913	7	3	7.913
2	0	3.783	7	5	11.016	6	5	11.016	2	0	2.833	7	4	8.825	7	4	8.825	2	0	2.395	7	4	6.207	7	4	6.207
2	1	4.725	7	6	11.773	6	6	11.773	2	1	3.411	7	5	9.365	7	5	9.365	2	1	2.887	7	5	8.391	7	5	8.391
2	2	5.633	8	0	7.916	6	7	7.916	2	2	4.405	8	0	6.550	8	0	6.550	2	2	3.830	8	0	5.891	8	0	5.891
2	3	6.496	8	1	8.410	6	8	8.410	2	3	5.173	8	1	7.809	8	1	7.809	2	3	4.366	8	1	6.279	8	1	6.279
2	4	7.141	8	2	9.312	6	9	9.312	2	4	5.930	8	2	8.653	8	2	8.653	2	4	5.260	8	2	7.215	8	2	7.215
2	5	8.145	8	3	10.090	6	10	10.090	2	5	6.653	8	3	9.510	8	3	9.510	2	5	5.940	8	3	7.736	8	3	7.736
2	6	8.936	8	4	10.866	6	11	10.866	2	6	7.372	8	4	10.286	8	4	10.286	2	6	6.622	8	4	8.547	8	4	8.547
2	7	9.712	8	5	11.613	6	12	11.613	2	7	8.080	8	5	11.055	8	5	11.055	2	7	7.295	8	5	8.943	8	5	8.943
2	8	10.476	9	0	8.530	6	13	8.530	2	8	8.589	8	6	12.349	8	6	12.349	2	8	7.961	9	0	6.432	9	0	6.432
2	9	11.021	9	1	9.011	6	14	9.011	2	9	9.472	9	0	13.129	9	0	13.129	2	9	8.622	9	1	6.814	9	1	6.814
2	10	11.759	9	2	9.902	6	15	9.902	2	10	9.969	9	1	14.001	9	1	14.001	2	10	9.277	9	2	7.611	9	2	7.611
3	0	4.520	9	3	10.672	6	16	10.672	3	0	3.611	9	2	14.886	9	2	14.886	3	0	3.097	9	3	8.267	9	3	8.267
3	1	5.375	9	4	11.443	6	17	11.443	3	1	4.067	9	3	15.771	9	3	15.771	3	1	3.511	9	4	9.057	9	4	9.057
3	2	6.247	10	0	9.137	6	18	9.137	3	2	4.975	10	0	16.656	10	0	16.656	3	2	4.651	10	0	6.971	10	0	6.971
3	3	7.334	10	1	9.607	6	19	9.607	3	3	5.724	10	1	17.541	10	1	17.541	3	3	5.074	10	1	7.345	10	1	7.345
3	4	7.918	10	2	10.490	6	20	10.490	3	4	6.325	10	2	18.426	10	2	18.426	3	4	5.628	10	2	8.139	10	2	8.139
3	5	8.700	10	3	11.251	6	21	11.251	3	5	7.183	10	3	19.311	10	3	19.311	3	5	6.314	10	3	8.789	10	3	8.789
3	6	9.480	11	0	9.737	6	22	9.737	3	6	7.895	10	4	20.196	10	4	20.196	3	6	6.990	11	0	7.505	11	0	7.505
3	7	10.246	11	1	10.197	6	23	10.197	3	7	8.795	11	0	21.081	11	0	21.081	3	7	7.659	11	1	7.874	11	1	7.874
3	8	11.229	11	2	11.074	6	24	11.074	3	8	9.293	11	1	21.966	11	1	21.966	3	8	8.322	11	2	8.665	11	2	8.665
4	0	5.222	11	3	11.826	6	25	11.826	4	0	10.174	11	2	22.851	11	2	22.851	4	0	8.982	11	3	9.310	11	3	9.310
4	1	6.028	12	0	10.331	6	26	10.331	4	1	4.211	12	0	23.736	12	0	23.736	4	0	9.666	12	0	8.036	12	0	8.036
4	2	6.878	12	1	11.783	6	27	11.783	4	2	4.689	12	1	24.621	12	1	24.621	4	1	10.099	12	1	8.407	12	1	8.407
4	3	7.703	12	2	11.654	6	28	11.654	4	3	5.555	12	2	25.506	12	2	25.506	4	2	10.985	12	2	9.188	12	2	9.188
4	4	8.518	13	0	10.921	6	29	10.921	4	4	6.478	13	0	26.391	13	0	26.391	4	3	11.871	13	0	8.564	13	0	8.564
4	5	9.293	13	1	11.365	6	30	11.365	4	5	7.037	13	1	27.276	13	1	27.276	4	4	12.756	13	1	8.924	13	1	8.924
4	6	10.065	14	0	11.506	6	31	11.506	4	6	7.735	14	0	28.161	14	0	28.161	4	5	13.641	14	0	9.089	14	0	9.089

A somewhat more complicated case making use of the possibility of combining component data when some sample sizes are equal, as described in Section 1, is illustrated by

Example 2. Consider a five-component series system with test data  $X_1 = 0, X_2 = 4, X_3 = 2, X_4 = 1, X_5 = 3$ , based on corresponding sample sizes  $n_1 = n_2 = n_3 = 300, n_4 = n_5 = 200$ . Here  $c = (3/300 + 2/200) = .02, a_1 = a_2 = a_3 = 1/300(.02) = 1.6, a_4 = a_5 = 1/200(.02) = 1/4$ . For the equivalent  $k^* = 2$  problem we divide the two distinct values of the  $a_i$ 's by their sum  $c_0 = 1/6 + 1/4 = 5/12$ , obtaining  $a_1^* = .40, a_2^* = .60$ . The corresponding numbers of failures are  $X_1^* = X_1 + X_2 + X_3 = 6$  and  $X_2^* = X_4 + X_5 = 4$ . For the reduced problem we consult Table 1 for  $\alpha = .10$  to find the 90 percent confidence bound 7.564. This must be multiplied by  $c_0$  to obtain the bound for the original  $\theta(\lambda)$ . The 90 percent lower bound  $r(x)$  for the system reliability given by (1.6) thus becomes  $1 - cc_0 t(x) = 1 - (.02)(5/12)(7.564) = .937$ .

If the observed numbers of failures fall outside the limits of Table 1, the approximate methods discussed in the next section may be employed.

#### 4. Approximations

In this section approximate confidence bounds for systems with more than two components having distinct sample sizes are developed. The use of the maximum likelihood ratio confidence bound for cases falling outside the scope of Table 1 is also discussed.

For  $k > 2$  it is impractical to generate the optimal ordering and the corresponding values of the upper confidence bound for more than a few illustrative cases. Thus, some method of approximating solutions for  $k > 2$  with acceptable precision is required. The approach to be followed here is to find a  $k = 2$  problem which is sufficiently similar in structure to the given  $k > 2$  problem so that the confidence bounds for the two problems are essentially the same except for a normalizing factor. This method may be thought of as an extension and refinement of the Lindstrom-Madden procedure (see Lloyd and Lipow 1977, and cf. Harris and Soms 1980).

The Lindstrom-Madden method first estimates the reliability by maximum likelihood and then uses the  $k = 1$  confidence bound solution for the component with the smallest sample size and a fictitious number of failures determined so as to reproduce the estimated system reliability. The procedure proposed here is to estimate two quantities, the value of  $\theta$  and the variance of the maximum likelihood estimate of  $\theta$ , and to use these estimates to find a  $k = 2$  problem based on two of the original  $a_i$ 's with a pair of corresponding fictitious observation values chosen to reproduce the estimated quantities. The two  $a_i$ 's are chosen to be as large as possible (corresponding to sample sizes as small as possible) subject to two constraints. The

first constraint guarantees that the resulting fictitious observations are non-negative. The second constraint requires that  $a_i$ 's corresponding to zero failures in the original problem not be considered unless all but one of the components have zero failures. These considerations lead to a unique  $k = 2$  problem whose solution provides a very good approximation to the solution for the original  $k > 2$  problem.

The elimination of  $a_i$ 's for components exhibiting zero failures is justified by the fact that the maximum likelihood ratio confidence bound (discussed later) is invariant under such transformations. That is, if the dimension  $k$  is reduced by the elimination of all  $a_i$ 's corresponding to zero failures, then the value of the maximum likelihood ratio bound remains unchanged.

The choice of the two quantities whose estimates are used to determine the pair of fictitious observations is supported by analogy with Lindstrom-Madden (in the case of  $\theta$ ) and by the fact that the two estimates are the ingredients of the asymptotic maximum likelihood ratio confidence bound thereby insuring asymptotic optimality. The details of the approximation algorithm are as follows:

(a) First  $\theta(\lambda) = \sum_{i=1}^k a_i \lambda_i$  is estimated by

$$\hat{\theta} = \sum_{i=1}^k a_i X_i, \quad (4.1)$$

and the quantity  $\text{Var}(\hat{\theta}) = \sum_{i=1}^k a_i^2 \lambda_i$  is estimated by

$$\hat{v} = \sum_{i=1}^k a_i^2 X_i. \quad (4.2)$$

(b) Next if at least one component exhibits one or more failures, the pair  $(a_i, a_j)$ ,  $i < j$ , is selected so that

$$(i) \quad a_i \leq \hat{v}/\hat{\theta} \quad , \quad \text{and}$$

$$(ii) \quad a_j \geq \hat{v}/\hat{\theta} \quad .$$

Subject to these conditions,  $a_i$  and  $a_j$  are taken to be the largest available values associated with at least one failure. If all  $a_i$ 's satisfying (i) correspond to zero failures, then  $a_i$  is taken to be the largest in that group. If all the  $a_j$ 's satisfying (ii) correspond to zero failures, then  $a_j$  is taken to be  $a_k$ . If all components exhibit zero failures,  $\hat{v}/\hat{\theta}$  is indeterminate and  $a_i$  and  $a_j$  are taken to be  $a_1$  and  $a_k$  respectively.<sup>1</sup>

(c) The pseudo-observations  $x_1^*$  and  $x_2^*$  are computed by the formulae

$$x_1^* = \frac{a_j \hat{\theta} - \hat{v}}{a_i (a_j - a_i)} \quad , \quad (4.3)$$

$$x_2^* = \frac{\hat{v} - a_i \hat{\theta}}{a_j (a_j - a_i)} \quad .$$

These values will be non-negative by conditions (i) and (ii) of (b), and when associated with  $a_i$  and  $a_j$  respectively they

---

<sup>1</sup>For this case the resulting confidence bound is exact and is the same as would be obtained by multiplying  $a_k$  times the upper confidence bound for a single Poisson parameter when zero failures are observed, i.e.,  $t(0) = a_k \log(1/\alpha)$ .

reproduce the values of  $\hat{\theta}$  and  $\hat{v}$  provided all other observations are replaced by zeros.

- (d) The  $k = 2$  problem with  $x_1^*$  and  $x_2^*$  associated with  $a_1^* = a_i/(a_i + a_j)$  and  $a_2^* = a_j/(a_i + a_j)$  respectively may now be treated using Table 1 to yield  $t(x_1^*, x_2^*)$ . Since  $x_1^*$  and  $x_2^*$  are not necessarily integers, it may be necessary to interpolate with respect to these arguments as well as the value of  $a_1 = a_1^*$ .

- (e) The approximate upper confidence bound  $t^*$  for  $\theta$  for the original  $k > 2$  problem is then given by

$$t^*(x) = (a_i + a_j)t(x_1^*, x_2^*) \quad . \quad (4.4)$$

In order to check the validity of this approximation algorithm, the confidence bounds for the first 24 points in the optimal (two-stage prospective) ordering were computed for a typical  $k = 3$  case. These results were obtained by a rather laborious method based on formula (2.7) and involving repeated interactive searches of the  $(\lambda_1, \lambda_2, \lambda_3)$  simplex. The approximation algorithm was applied to each of these points and the comparative results are shown in Table 2. The values of  $a_1$ ,  $a_2$ , and  $a_3$  for this example were chosen to be of roughly the same magnitude but not so close that the combination of any two would be indicated.

To check the algorithm for cases farther from the origin several additional examples were considered using a somewhat different method which avoids the necessity for sequentially generating the

Table 2. The Performance of the Approximation Algorithm for the First 24 Ordered Points for a Typical Example With  $k = 3$ .

$\alpha = .05; a_1 = .2, a_2 = .3, a_3 = .5$						
Two-Step Prospective Sequential					Approximation Algorithm	Relative Error
n	$x_1$	$x_2$	$x_3$	$t(x)$	$t^*(x)$	
1	0	0	0	1.498	1.498	+ 00.0 %
2	1	0	0	1.553	1.555	+ 00.1
3	0	1	0	1.656	1.663	+ 00.4
4	2	0	0	1.703	1.705	+ 00.2
5	1	1	0	1.756	1.540	- 12.3
6	2	1	0	1.861	1.752	- 05.9
7	3	0	0	1.933	1.891	- 02.1
8	0	2	0	1.981	1.995	+ 00.7
9	1	2	0	2.085	2.052	- 01.6
10	4	0	0	2.088	2.094	+ 00.3
11	3	1	0	2.162	1.991	- 07.9
12	2	2	0	2.268	2.208	- 02.7
13	5	0	0	2.300	2.309	+ 00.4
14	0	3	0	2.354	2.397	+ 01.8
15	0	0	1	2.372	2.372	+ 00.0
16	4	1	0	2.382	2.291	- 03.9
17	1	0	1	2.429	2.431	+ 00.1
18	1	3	0	2.472	2.438	- 01.4
19	5	1	0	2.504	2.486	- 00.7
20	0	1	1	2.543	2.529	- 00.5
21	6	0	0	2.575	2.551	- 01.0
22	2	0	1	2.602	2.589	- 00.8
23	2	3	0	2.609	2.641	+ 01.2
24	1	1	1	2.660	2.668	+ 00.3

optimal ordering of the sample points. For these examples the ordering was generated by the values of the maximum likelihood ratio confidence bounds (see below) associated with the sample points. Since this ordering is asymptotically optimal (for large  $\lambda_i$ 's), it can be expected to produce good results for sample points well removed from the origin. The values of the confidence bounds calculated using (2.7) and the approximations obtained by the proposed algorithm are shown for these examples in Table 3.

In Table 3 the tendency of the computed values of  $t(x)$  to be slightly larger than the algorithm values may be due to the fact that the ordering used to compute the former is non-optimal. Formal application of the algorithm may occasionally result in the selection of nearly equal values  $a_i$  and  $a_j$  in step (b). When this happens, improved results may be obtained by first reducing the dimension  $k$  by combining the nearly equal  $a_i$ 's and then applying the algorithm. This method was used for the two cases in Table 3 marked by (+). The algorithm values for all cases where  $a = (.14, .16, .70)$  or  $a = (.15, .41, .44)$  do not differ substantially from the values which would be obtained by reducing to the  $k = 2$  case by combining the nearly equal  $a_i$ 's. Similarly, the values for the cases where  $a = (.32, .33, .35)$  can be nearly reproduced by multiplying the  $k = 1$  bound by  $\bar{a} = .333$ .

The number of sample points appearing in the ordering before a given level of  $t(x)$  is reached increases rapidly as  $k$  increases.



Table 3. Several Examples Comparing the Algorithm Values With the Exact Bounds Based on the Ordering Generated by the m&rb.

$\alpha = .05$									
$a_1$	$a_2$	$a_3$	$x_1$	$x_2$	$x_3$	Position in Ordering	$t(x)$	Algorithm Value $t^*(x)$	m&rb
.20, .30, .50			2, 2, 1			50	3.329	3.273	3.070
			5, 0, 2			100	4.068	3.957	3.798
			6, 6, 0			200	4.887	4.789	4.708
			9, 1, 3			400	5.980	5.892	5.795
.14, .16, .70			5, 4, 0			50	3.026	3.084 <sup>†</sup>	2.218
			2, 10, 0			100	3.424	3.531 <sup>†</sup>	2.921
			2, 5, 1			200	4.407	4.086	3.729
.15, .41, .44			13, 0, 0			50	3.245	3.223	2.980
			5, 2, 1			100	4.028	3.860	3.699
			3, 3, 2			200	4.861	4.738	4.559
.32, .33, .35			0, 3, 2			50	3.492	3.518	3.256
			1, 5, 1			100	4.339	4.332	4.076
			1, 5, 3			200	5.198	5.234	4.993

<sup>†</sup> Reduction to  $k=2$  case by combining  $a_1$  and  $a_2$ .

Hence for  $k > 2$  the algorithm may be applied to sample points positioned quite far along in the ordering without requiring values beyond the scope of Table 1, as is seen in the examples of Table 3.

The results detailed in Tables 2 and 3 suggest that the application of the proposed algorithm together with the combining of nearly equal  $a_i$ 's when indicated will nearly always produce confidence bounds for  $\theta(\lambda)$  subject to relative errors not exceeding 10 percent and often much less. Furthermore, the lower confidence bound for reliability  $r(x)$  given by (1.6) will exhibit a much smaller relative error since the relative error in approximations for  $t(x)$  applies only to the difference between the lower bound and one; a quantity which at worst is of the order of  $1/10$  in the contemplated applications.

In situations where very large sample sizes are available, it may happen that the observed numbers of failures exceed the limits of Table 1 even though the system reliability is high. For such cases the maximum likelihood ratio bound (mlrb) may be used. This approximate confidence bound is obtained in the usual way from the maximum likelihood ratio statistic for testing the hypothesis  $H_0: \theta(\lambda) = \theta_0$  versus all alternatives. The corresponding one-sided confidence bound may be shown (see Johns 1975) to be determined as follows: Let  $\hat{\mu}$  be the positive real root less than  $1/(\text{largest } a_i \text{ for which } X_i > 0)$  of the equation

$$\chi_{1,2\alpha}^2 = 2 \sum_{i=1}^k X_i \left\{ \frac{a_i \hat{\mu}}{1 - a_i \hat{\mu}} + \log(1 - a_i \hat{\mu}) \right\}, \quad (4.5)$$

where  $\chi_{1,2\alpha}^2$  is the upper  $100(2\alpha)$ -th percentage point of the chi-squared distribution with one degree of freedom. Then the quantity

$$\hat{t}(X) = \sum_{i=1}^k a_i X_i / (1 - a_i \hat{\mu}) \quad (4.6)$$

is the approximate upper  $1 - \alpha$  level confidence bound for  $\theta(\underline{\lambda})$ . As  $\max(\lambda_1, \lambda_2, \dots, \lambda_n) \rightarrow \infty$ , the mlrb  $\hat{t}(X)$  may be shown (see Johns 1975) to be asymptotically equivalent to

$$\tilde{t}(X) = \sum_{i=1}^k a_i X_i + z_{\alpha} \left( \sum_{i=1}^k a_i^2 X_i \right)^{1/2}, \quad (4.7)$$

where  $z_{\alpha}$  is the  $100\alpha$ -th percentage point of the standard normal distribution.

Neither of these approximate bounds is useful for sample points near the origin in the usual orderings, but  $\hat{t}$  given by (4.6) becomes sufficiently precise for application to sample points beyond the scope of Table 1. A comparison of  $t(x_1, x_2)$  and the corresponding mlrb for the last (i.e., the 100-th) points in each of the optimal orderings for the cases covered in Table 1 is given in Table 4.

Table 4.  $m\ellrb/t(x_1, x_2)$  for the 100-th Point in Each Ordering

$\alpha$	$a_1$								
	.05	.10	.15	.20	.25	.30	.35	.40	.45
.01	.62	.95	.97	.97	.98	.97	.98	.97	.99
.05	.72	.88	.96	.95	.98	.97	.99	.95	.96
.10	.89	.95	.95	.96	.97	1.00	.97	.96	.98

These results suggest that the  $m\ellrb$  possesses satisfactory precision for sample points beyond those listed in Table 1 for  $k = 2$  whenever  $a_1 \geq .10$ . The last column of Table 3 giving the  $m\ellrb$  values for the examples considered illustrates the fact that for  $k > 2$  the  $m\ellrb$  tends to underestimate the correct value of the bound for sample points within the range that can be dealt with using Table 1 and the algorithm.

Two potential sources of error for the lower confidence bound on system reliability remain to be discussed. They are (i) the Poisson approximation to the binomial distribution of the observed failures, and (ii) the approximation for reliability given in (1.2) and reflected in the formula (1.6) for the lower bound  $r(x)$ . It is intuitively clear from (1.1) ff. that the "worst case" for the Poisson approximation should occur when  $k = 1$ ; since to match a given  $k = 1$  level of reliability, say  $1 - p$ , by a  $k > 1$  case, we must have  $p \approx \sum_{i=1}^k p_i$ , so that the  $p_i$ 's must be smaller than  $p$  which tends to improve the Poisson approximation.

For the case  $k = 1$  the familiar upper confidence bounds for a single Poisson parameter apply and the actual coverage probabilities for the proposed method (3.1) can be computed for any  $n$  and  $p$  from tables of the binomial distribution. The results of several such calculations are shown in Table 5.

Table 5. "Worst Case" Analysis ( $k=1$ ). Minimum Coverage Probabilities for  $r(x)$ .

Reliability $q = 1 - p$	$1 - \alpha$		
	.90	.95	.99
.95	.906	.954	.991
.90	.912	.958	.992
.80	.924	.965	.995
.70	.936	.972	.997

These values suggest that the approximations operate to make the proposed confidence bounds slightly conservative. It is interesting to observe that the minimum coverage probabilities are not drastically different from the nominal values, even for a true reliability as low as .70.

##### 5. Acknowledgments and Remarks Concerning the Computations

The computations for Table 1 were performed on a Digital Equipment PDP-11/34 running under a UNIX operating system. The production program for these computations (506 lines) was written in the "C" language by the author. The method used for the

computation of the upper bounds was an implementation of (2.7); and as has been noted, the sample points were ordered by the two-step prospective sequential method. The tabled results were subjected to various checks to insure that the correct global maxima were found in each case.

The computations involved in obtaining the one-step look-ahead results and the tree analysis associated with the semi-Bayesian results discussed in Section 3 were done on an IBM 370/168 machine using Fortran programs developed by David Pasta. These programs compute the upper bounds by a somewhat more complicated method used in the earlier phases of the study and detailed in Johns (1975).

Thanks are also due to Barry Eynon who helped with the development of Figure 1 and the display of Table 1, and to Robert Bell and Keaven Anderson for careful readings of Sections 2.

## REFERENCES

- BOL'SHEV, L.N., and LOGINOV, E.A. (1969), "Interval Estimates in the Presence of Noise," Theory of Probability and Its Applications, 11, 82-94.
- BUEHLER, R.J. (1957), "Confidence Intervals for the Product of Two Binomial Parameters," Journal of the American Statistical Association, 52, 482-493.
- BUTCHER, A., LAMPKIN, H., and WINTERBOTTOM, A. (1978), "Transformations Improving Maximum Likelihood Confidence Intervals for System Reliability," Technometrics, 20, 467-473.
- EL MAWAZINY, A.H., and BUEHLER, R.J. (1967), "Confidence Limits for the Reliability of a Series Systems," Journal of the American Statistical Association, 62, 1452-1459.
- HARRIS, B., and SOMS, A.P. (1980), "Bounds for Optimal Confidence Limits for Series Systems," Technical Summary Report No. 2093, University of Wisconsin-Madison, Mathematics Research Center.
- \_\_\_\_\_. (1981), "Improved Sudakov-Type Bounds for Optimal Confidence Limits on the Reliability of Series Systems," Technical Report No. 637, University of Wisconsin-Madison, Department of Statistics.
- JOHNS, M.V., JR. (1975), "Reliability Assessment for Highly Reliable Systems," Technical Report No. 1, Stanford University, Department of Statistics.
- JOHNS, M.V., JR. (1977), "Some Aspects of Reliability Assessment Based on Sample Orderings," Proceedings of the ARO Workshop on Reliability and Probabilistic Design, 101-110.
- JOHNSON, J.R. (1969), "Confidence Interval Estimation of the Reliability of Multicomponent Systems Using Component Test Data," PhD dissertation, University of Delaware.
- LEHMANN, E.L. (1959), Testing Statistical Hypotheses, New York: John Wiley and Sons.
- LENTER, M.M., and BUEHLER, R.J. (1963), "Some Inferences About Gamma Parameters with an Application to a Reliability Problem," Journal of the American Statistical Association, 58, 670-677.
- LIEBERMAN, G.J., and ROSS, S.M. (1971), "Confidence Intervals for Independent Exponential Series Systems," Journal of the American Statistical Association, 66, 837-840.

- LIPOW, M., and RILEY, J. (1959), "Tables of Upper Confidence Limits on Failure Probability of 1, 2, and 3 Component Serial Systems," Report No. TR-59-0000-00756, Space Technology Laboratories, Inc.
- LLOYD, D.K., and LIPOW, M. (1977), Reliability: Management, Methods, and Mathematics, 2nd ed., Redondo Beach: Published by the Authors.
- MADANSKY, A. (1965), "Approximate Confidence Limits for the Reliability of Series and Parallel Systems," Technometrics, 7, 495-503.
- MANN, N.R., and GRUBBS, F.E. (1974), "Approximately Optimum Confidence Bounds for System Reliability Based on Component Test Data," Technometrics, 16, 335-348.
- \_\_\_\_\_, SCHAFER, R.E., and SINGPURWALLA, N.D. (1974), Methods for Statistical Analysis of Reliability and Life Data, New York: John Wiley and Sons.
- PEARSON, E.S., and HARTLEY, H.O. (1958), Biometrika Tables for Statisticians, Vol. I, 2nd Ed., Cambridge: University Press.
- STECK, G.P. (1957), "Upper Confidence Limits for the Failure Probability of Complex Networks," Technical Report No. SC-433, Sandia Corporation.
- WINTERBOTTOM, A. (1974), "Lower Limits for Series System Reliability from Binomial Data," Journal of the American Statistical Association, 69, 782-788.
- WINTERBOTTOM, A. (1980), "Asymptotic Expansions to Improve Large Sample Confidence Intervals for System Reliability," Biometrika, 67, 351-357.



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER No. 6	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  FEASIBLE OBJECTIVE CONFIDENCE BOUNDS FOR SYSTEM RELIABILITY		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  M. Vernon Johns, Jr.		8. CONTRACT OR GRANT NUMBER(s)  ARO DAAG29-79-C-0166
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, California		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709		12. REPORT DATE August 20, 1981
		13. NUMBER OF PAGES 45
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Reliability, series-system, confidence bounds, Poisson approximation,  Lindstrom-Madden, sample orderings		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The basic problem of determining objective (frequentistic) confidence bounds for the reliability of a series system based on failure data from tests of the independent components is addressed. The notion of confidence bounds based on orderings imposed on the sample space is exploited, and certain optimality considerations are incorporated. Advantage is taken of the simplifications resulting from the use of the Poisson approximation for data from highly reliable components. Tables of exact confidence bounds are produced for the case of two-component systems. These bounds are computed using sample orderings		

generated sequentially by a two-stage, prospective optimization procedure. A generalization of the Lindstrom-Madden technique is proposed for using the tables to find confidence bounds for systems consisting of more than two components with differing sample sizes.